



LJMU Research Online

King, NG, Wilcockson, DC, Webster, R, Smale, DA, Hoelters, LS and Moore, PJ

Cumulative stress restricts niche filling potential of habitat-forming kelps in a future climate

<http://researchonline.ljmu.ac.uk/id/eprint/11450/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

King, NG, Wilcockson, DC, Webster, R, Smale, DA, Hoelters, LS and Moore, PJ (2017) Cumulative stress restricts niche filling potential of habitat-forming kelps in a future climate. Functional Ecology, 32 (2). pp. 288-299. ISSN 0269-8463

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

Cumulative stress restricts niche filling potential of habitat-forming kelps in a future climate

Nathan G. King¹  | David C. Wilcockson¹ | Richard Webster¹ |
Dan A. Smale²  | Laura S. Hoelters¹ | Pippa J. Moore^{1,3}

¹Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Aberystwyth, UK

²Marine Biological Association of the United Kingdom, The Laboratory, Plymouth, UK

³Centre for Marine Ecosystems Research, School of Natural Sciences, Edith Cowan University, Joondalup, WA, Australia

Correspondence

Nathan G. King
Email: nak14@aber.ac.uk

Funding information

Institute of Biological, Environmental and Rural Sciences PhD studentship; Marie Curie Career Integration Grant, Grant/Award Number: PCIG10-GA-2011-303685; Aberystwyth University Research Fund, Grant/Award Number: 1211Z-01; Natural Environment Research Council of the UK, Grant/Award Number: NE/K008439/1

Handling Editor: Emily Carrington

Abstract

1. Climate change is driving range contractions and local population extinctions across the globe. When this affects ecosystem engineers the vacant niches left behind are likely to alter the wider ecosystem unless a similar species can fulfil them.
2. Here, we explore the stress physiology of two coexisting kelps undergoing opposing range shifts in the Northeast Atlantic and discuss what differences in stress physiology may mean for future niche filling.
3. We used chlorophyll fluorescence (F_v/F_m) and differentiation of the heat shock response (HSR) to determine the capacity of the expanding kelp, *Laminaria ochroleuca*, to move into the higher shore position of the retreating kelp, *Laminaria digitata*. We applied both single and consecutive exposures to immersed and emersed high and low temperature treatments, replicating low tide exposures experienced in summer and winter.
4. No interspecific differences in HSR were observed which was surprising given the species' different biogeographic distributions. However, chlorophyll fluorescence revealed clear differences between species with *L. ochroleuca* better equipped to tolerate high immersed temperatures but showed little capacity to tolerate frosts or high emersion temperatures.
5. Many patterns observed were only apparent after consecutive exposures. Such cumulative effects have largely been overlooked in tolerance experiments on intertidal organisms despite being more representative of the stress experienced in natural habitats. We therefore suggest future experiments incorporate consecutive stress into their design.
6. Climate change is predicted to result in fewer ground frosts and increased summer temperatures. Therefore, *L. ochroleuca* may be released from its summer cold limit in winter but still be prevented from moving up the shore due to desiccation in the summer. *Laminaria ochroleuca* will, however, likely be able to move into tidal pools. Therefore, only partial niche filling by *L. ochroleuca* will be possible in this system as climate change advances.

KEYWORDS

climate change, heat shock response, *Laminaria digitata*, *Laminaria ochroleuca*, leading edge, range shift, Trailing edge

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2017 The Authors. *Functional Ecology* published by John Wiley & Sons Ltd on behalf of British Ecological Society.

1 | INTRODUCTION

The world's oceans have warmed by 0.11°C per decade for the last 40 years while short term rapid increases in ocean and aerial temperatures (heatwaves) have also increased in frequency and duration (Christidis, Stott, Brown, Hegerl, & Caesar, 2005; Coumou & Rahmstorf, 2012; Hartmann, Tank, & Rusticucci, 2013; Lima & Wethey, 2012; Meehl & Tebaldi, 2004; Seneviratne et al., 2012). Together, these two types of warming are resulting in the global redistribution of marine species (Burrows et al., 2011; Hobday et al., 2016; Poloczanska et al., 2013; Smale & Wernberg, 2013), altering the structure and functioning of entire ecosystems (Doney et al., 2012; Hawkins et al., 2009; Wernberg et al., 2016). In marine systems, organisms generally occupy the entirety of their "thermal niche" meaning marine populations are particularly responsive to warming (Sunday, Bates, & Dulvy, 2012). While local conditions can cause mosaic patterns of stress intensities in the intertidal (Helmuth et al., 2016; Lourenço et al., 2016), it is still range margins that are seen to be at the forefront of warming trends, resulting in poleward range expansions (leading edge) or contractions (trailing edge) (Chen, Hill, Ohlemüller, Roy, & Thomas, 2011; Hampe & Petit, 2005; Poloczanska et al., 2013). Understanding the physical determinants that set these boundaries is therefore critical for predicting the structure and functioning of ecosystems as climate change progresses (Gaston, 2003).

Kelps dominate shallow temperate rocky reefs where they support high levels of primary productivity and modify environmental conditions such as light (Wernberg, Kendrick, & Toohey, 2005), water flow (Rosman, Koseff, Monismith, & Grover, 2007), physical disturbance (Connell, 2003) and sedimentation rates (Eckman, Duggins, & Sewell, 1989), allowing rich assemblages to persist (Steneck et al., 2002). The biogeographic ranges of kelp species are largely controlled by temperature (Eggert, 2012; Lüning & Tom Dieck, 1990). As such, increases in mean temperatures and heatwaves can alter macroalgal distributions (Edwards & Estes, 2006; Filbee-Dexter, Feehan, & Scheibling, 2016; Smale & Wernberg, 2013; Smale, Wernberg, Yunnice, & Vance, 2015; Tuya et al., 2012; Voerman, Llera, & Rico, 2013), which can have catastrophic consequences for the associated ecosystems they underpin (Wernberg et al., 2013, 2016).

The British Isles is an important Northeast Atlantic biogeographical transition zone (Forbes, 1853). Along Britain's south coast, two structurally similar canopy-forming kelps occur in sympatry at opposing edges of their distributional ranges. *Laminaria digitata*, a boreal species, approaches its trailing edge while *Laminaria ochroleuca*, a Lusitanian species, reaches its leading edge (Bartsch, Vogt, Pehlke, & Hanelt, 2013; Smale et al., 2015). Both coexist on moderately sheltered to moderately exposed shores where their vertical distributions overlap. *Laminaria digitata* dominates the low intertidal (<1 m above Chart datum) before being replaced by *L. ochroleuca* further down the shore (Hargrave, Foggo, Pessarrodona, & Smale, 2016). These species are currently undergoing opposing poleward range migrations, with *L. digitata* population declines observed along both sides of the English Channel, while *L. ochroleuca* has proliferated at its leading range edge (Raybaud et al., 2013; Smale et al., 2015). *Laminaria digitata* is a key

ecosystem engineer (Schultze, Janke, Krüß, & Weidemann, 1990) and its loss could significantly impact wider community structure, unless another functionally similar species, such as *L. ochroleuca*, is able to move into the vacated habitat left behind.

The environmental conditions a species can survive in (fundamental niche) are often wider than where it is found (realised niche). The reason for this disparity is often due to interspecific competitive exclusion (Hutchinson, 1957, 1978). It is not known to what extent *L. ochroleuca*'s vertical distribution on the shore is dictated by an inability to tolerate greater periods of low tide stress or through *L. digitata*'s competitive dominance at higher tidal heights. Indeed, *L. ochroleuca* is found in much more varied habitats at lower latitudes that are currently occupied by *L. digitata* in Britain (e.g. tide pools & exposed coasts) (Pereira, Engelen, Pearson, Valero, & Serrão, 2015). The acquisition of such knowledge will allow for more accurate predictions of the structure and functioning of these communities in a future warmer world.

Optimal growth and maximal survival temperatures of macroalgae are well studied (Bolton & Anderson, 1987; Bolton & Lüning, 1982; Hargrave et al., 2016; Lüning, 1984; Lüning & Freshwater, 1988; Orfanidis, 1991; Simonson, Metaxas, & Scheibling, 2015; Tom Dieck, 1992) but the effects of consecutive low tide exposures remain relatively unknown (but see Pereira et al., 2015). This is surprising as low tide stress has long been known to be important in determining the vertical and latitudinal distributions of intertidal organisms (Dring, 1982; Evans, 1948) and increases in aerial temperatures can cause shifts in macroalgal distributions (Harley & Paine, 2009; Harley et al., 2012; Martínez et al., 2012; Ugarte, Critchley, Serdynska, & Deveau, 2009). Here we examined the potential for *L. ochroleuca* to move into niches left behind by *L. digitata* as its biogeographic range contracts in response to warming. Firstly, we determined the thermal tolerance of both species by subjecting individuals to a single temperature shock using a traditional metric of stress, upregulation of *Hsp70* across a range of temperatures. Secondly, we used chlorophyll fluorescence (F_v/F_m) to investigate the resistance and resilience of each species to consecutive low tide scenarios. Understanding tolerances to consecutive low tide stress of these two species will provide important insight into how climate change can alter range edge dynamics whilst also providing foresight into the future species composition of kelp-dominated communities on Northeast Atlantic rocky shores.

2 | MATERIALS AND METHODS

Mature *L. digitata* and *L. ochroleuca* sporophytes ($n = 5$) were collected on a low spring tide at St Mawes Bay, Cornwall, UK (50°9'24.4764" N, 5°1'4.1628" W) in April and September 2015. Heat shock experiments (qPCR) were conducted in April and consecutive stress assays were conducted in September. All individuals were taken from the overlapping zone where both species directly coexist (c. 0.4 m above chart datum). Individuals were transported to the laboratory in cool dark containers and held in aerated recirculating tanks for c. 7 days under photosynthetic flux density of c. 20 $\mu\text{mol m}^{-2} \text{s}^{-1}$

(12:12 hr). Holding tanks were maintained at ambient sea temperatures, 9°C (April) and 15°C (September). Following well-established methods in seaweed photophysiology and gene expression studies (García-Mendoza & Colombo-Pallotta, 2007; Henkel & Hofmann, 2008; Jueterbock et al., 2014; Pearson, Lago-Leston, & Mota, 2009; Rothäusler, Gómez, Karsten, Tala, & Thiel, 2011), we excised discs of tissue (diameter: 27 mm, area: 11.45 mm²) from each kelp, using a cork borer with discs left for 1 hr before being used in stress assays. To establish that excised discs responded similar to whole kelps, we measured F_v/F_m values (a measure of stress see below), which were similar across tissue types.

2.1 | Heat shock response

When environmental stresses cause proteins to denature and aggregate, organisms upregulate a suite of chaperone proteins known as Heat Shock Proteins (Hsps). These Hsps preserve normal cell function by ensuring appropriate protein folding during translation (Frydman, 2001), membrane stability, and transport (Hartl & Hayer-Hartl, 2002) and protein refolding (Hendrick & Hartl, 1993). The level of Hsp transcript is a proxy for cellular damage and quantifying it across a range of experimental treatments allows powerful inferences regarding stress tolerance to be made. The heat shock response (HSR) is the induction of Hsps in response to temperature shocks (Parsell & Lindquist, 1993). Induction follows a bell shaped curve and specific thermal set points along this curve (T_{peak} and T_{off}) indicate different important aspects of thermotolerance (Tomanek & Somero, 1999, 2002). T_{peak} represents an organism's maximum thermal tolerance as past this point protein synthesis can no longer keep up with the increasing thermal insult. T_{off} is an ultimate upper limit in the functioning of the translational machinery (Barua & Heckathorn, 2004; Tomanek, 2010). Both T_{peak} and T_{off} are fixed set points of the HSR and are not influenced by previous thermal history (Barua & Heckathorn, 2004). Therefore, seasonal differences in sampling periods between HSR experiments (spring) and chlorophyll fluorescence (summer) will not affect methodological comparisons.

To compare the HSR of *L. digitata* and *L. ochroleuca* $n = 5$ kelp discs per treatment were heat shocked at one of seven temperatures (8°C, 12°C, 16°C, 20°C, 24°C, 28°C or 32°C) for 1 hr. Discs were held in individual 100 ml sample pots in thermostatically controlled 35 L water baths with recirculating, aerated seawater. Discs were then removed, blotted dry and snap frozen in liquid N₂ and stored at -80°C until RNA extraction.

2.2 | RNA extraction and qPCR

Total RNA was extracted following a protocol from (Pearson, Lago-Leston, Valente, & Serrão, 2006). All glassware and consumables were treated with RNaseAway (Ambion, UK) and rinsed with DEPC-treated water before use. All reagents were molecular grade and RNase free. Half a disc (c. 300 mg) of heat shocked kelp tissue was ground to a fine powder under liquid N₂ using a pestle and mortar, placed in an extraction buffer (100 mM Tris, 50 mM EDTA, 2 M NaCl and 2% CTAB

pH 7.5), vortexed and left at room temperature (RT) for 10–15 min. To minimise endogenous RNase activity, DTT was added (from 1 M stock) to the extraction buffer immediately prior to use to a final concentration of 50 mM. Total RNA was extracted from the resultant homogenate by chloroform extraction. One ml of chloroform:isoamyl alcohol (24:1 v/v) was added, vortexed vigorously, centrifuged at 12,000 g for 20 min at 20°C and the upper aqueous phase transferred to a new tube. Absolute EtOH was added (0.3 volumes) and gently mixed by rocking. Another chloroform extraction was then performed on this upper aqueous phase/EtOH mixture under the same conditions as the first. RNA was precipitated with 0.25 vol 12 M LiCl in the presence of 1% mercaptoethanol at -80°C for 30 min. Samples were centrifuged at 14,000 g for 30 min at 4°C and the pellet washed with 70% EtOH and recollected by centrifugation at 14,000 g for 10 min at RT. The RNA pellet was air-dried and then suspended in 50 µl DEPC-treated water. Contaminating gDNA was digested, using 2U DNase 1 (Turbo-DNase free) (Ambion, UK) following the manufacturer's instructions and stored at -80°C. Total RNA (100 ng) was reversed transcribed using High Capacity cDNA Reverse Transcription Kit (Applied Biosystems) according to manufacturer's instructions and using random hexamer primers.

In order to determine the sequence of the *Hsp70* gene for *L. ochroleuca* and *L. digitata*, PCR primers were designed based on *Saccharina japonica* (accession no FJ375359.1). PCRs were run in 20 µl volumes, using Bioline MyTaq Red PCR mix with 100 µM F and R primers and 1 µl cDNA template. Cycling parameters were 5 min at 95°C, followed by 40 cycles of 15 s at 95°C, 60 s at 55°C and 60 s at 72°C and a final extension of 10 min at 72°C. Amplicons were resolved on 1.5% agarose gels and products of the expected size were excised and extracted, using Bioline Isolate II clean-up columns. All products were subjected to direct, Sanger sequencing in-house.

PRIMER 3 software (Life Technologies) was used to design all qPCR primers. *Hsp70* primers were based on the consensus sequences for *L. ochroleuca* and *L. digitata*. For *L. digitata* gene expression normalisation, two reference genes were used, 18s ribosomal RNA (accession no. JX442505) and Rubisco large subunit (accession no. AY157696). For *L. ochroleuca* expression normalisation only used 18s Ribosomal RNA (Table 1).

Quantitative PCR reactions were run in triplicate on 384 well plates on an Applied Biosystem Quant Studio 120-Flex platform, with each plate containing all targets and housekeeping genes for each species. Each well contained 1 µl of cDNA in 10 µl SYBR green reactions (Bioline Sensifast Lo-Rox kit). The PCR amplification protocol consisted of 95°C for 2 min followed by 40 cycles of 95°C for 5 s and 60°C for 15 s (for all assays). Amplification specificity was verified by melt curve analysis (from 60°C to 95°C) and agarose gel electrophoresis. Four 10-fold serial dilutions of cDNA product were used to verify that all amplification reactions had efficiencies in the range of 90%–105%. No reverse transcriptase and no template controls were performed for each assay plate to ensure no contamination from gDNA was present. Relative mRNA levels were calculated as follows: Firstly, the difference in Ct values from *Hsp70* and the internal reference gene were calculated (Δ CT). The Δ CT value was then subtracted by the Δ CT of

TABLE 1 qPCR primer sequences used for assays on heat shocked *Laminaria digitata* and *Laminaria ochroleuca*

	Forward	Reverse
<i>L. digitata</i>		
<i>Hsp70</i>	GCTGCGAGTCGTTGAAGTA	TGGTGCTCGTGAAGATGAAG
<i>18s rRNA</i>	CGGAAGGATCATTACCGAAA	CCCAACTTCGCATAACGAAT
<i>Rubisco</i>	GACATGGATTGGGCATCTCTT	GTAGAACCACATCGTCACCTA
<i>L. ochroleuca</i>		
<i>Hsp70</i>	GAAGTTCGCTTGAAGTCTTG	GACGATCGAGGAGGGTATCT
<i>18s rRNA</i>	GATGAAGAACGAGCGAAATG	GTCAACAGACTCCGACAA

the control (8°C) from each individual ($\Delta\Delta CT$). Relative expression of *Hsp70* was then calculated by $e^{(-\Delta\Delta CT)}$.

2.3 | Consecutive stress experiments

The experimental design for consecutive stress assays was based on field observations. Both species are found on the low shore and are exposed to low tide stress together only on spring tides. Aerial exposure usually last c. 1 hr over each low tide period for four days before becoming immersed until the next spring tidal cycle.

Consecutive stress assays ($n = 5$) were conducted on two discs of tissue a mean value of which was used in all further analysis. Individual discs were placed under one of a number of different conditions for 1 hr, to simulate the low tide and then placed back into the control tanks (15°C) to simulate the returning tide. For immersed treatments, discs were held in individual 100 ml sample pots in thermostatically controlled 35 L water baths with recirculating, aerated seawater. After 24 hrs recovery in control tanks, discs were again exposed to their respective 1 hr treatments. In total, four simulated tidal cycles were performed on the same disc. Recovery was subsequently monitored daily for three days. Stress measurements (F_v/F_m see below) were taken directly before and after each “low tide” treatment.

Treatments consisted of either emerged or immersed conditions and tested tolerance to either high or low temperatures. Conditions were: control at 15°C, immersion at 24°C, 28°C, 32°C (simulating summer tidal pools), 0°C and 4°C (simulating winter tidal pools), emersion at 24°C, 28°C, and 32°C (simulating summer low tide aerial exposure) and -5°C (simulating severe winter ground frost). These temperature ranges were chosen to represent realistic field exposures. Low shore tidal pools in the study area can reach 28°C in summer (Martins, Hawkins, Thompson, & Jenkins, 2007) while blades of *L. ochroleuca* can be subjected to temperatures of >30°C in tidal pools in its southern range (Engelen et al., 2008). Similarly aerial temperatures can already reach 32°C in summer and ground frosts are encountered during winter spring tides (Moore, unpub).

The experimental design allows clear interspecific comparisons to be made for different types of immersion and emersion stress, yet some caution should be taken when generalising to field conditions. For example, only temperature was elevated in warm-emersed treatments whereas in natural settings a number of environmental factors contribute to low tide stress including relative humidity, wind speed and solar radiation. Moreover, the abrupt temperature shocks used

in this experiment are realistic for emersion stresses but are more rapid than the gradual warming seen in tidal pools. However, such shocks still allow direct comparisons of interspecific thermal tolerance (Henkel & Hofmann, 2008; Jueterbock et al., 2014; Smolina, Kollias, Jueterbock, Coyer, & Hoarau, 2016).

Stress was measured, using pulse amplitude modulated (PAM) fluorometry (Walz Diving PAM, Germany). The maximum quantum yield of PSII (ratio of photochemical quenching (F_v) to total fluorescence from closed PSII reaction centres (F_m)) is proportional to its efficiency (Butler, 1978) and F_v/F_m is an established method to quantify temperature stress in seaweeds (Jueterbock et al., 2014; Pearson et al., 2009; Saada et al., 2016). All F_v/F_m values were dark adapted for 15 min. The results for each species were normalised to one using the initial F_v/F_m values taken prior to the start of any treatments. This allowed small intrinsic interspecific differences in starting F_v/F_m to be accounted for.

For high temperature emersion treatments the level of desiccation experienced at each time point was expressed as relative water content (RWC %). Initial wet weights of fully hydrated discs were obtained at the start of the experiment prior to any experimental treatments (FW) and subtracted from their dry weights (DW) (estimated based on wet: dry weight of discs of both kelps, $n = 12$ for both *L. digitata* and *L. ochroleuca*, taken independently and dried at 60°C for 48 hrs). Wet weight of kelp discs at each experimental time point (IW) (pre- and post-treatments) was subtracted from the dry weight and then divided by FW-DW. The resulting value was multiplied by 100.

$$RWC = [(IW - DW)/(FW - DW)] \times 100$$

IW is wet weight of treated kelp; FW is initial wet weight of fully hydrated kelp (pre-treatments); DW is dry weight of kelp disc.

2.4 | Statistical analysis

Differences in upregulation of *Hsp70* to common garden temperature stress between species was measured, using univariate permutational ANOVA and the PERMANOVA module (Anderson, 2001) within PRIMER 6 software (Clarke & Gorley, 2006). The model included the two fixed factors; species (2 levels: *L. digitata*, *L. ochroleuca*) and temperature (7 levels: 8°C, 12°C, 16°C, 20°C, 24°C, 28°C & 32°C). Permutations (9,999 under a reduced model) were conducted on a similarity matrix constructed from Euclidean distances between untransformed data (relative *Hsp70* expressions). When conducting

PERMANOVA analysis on univariate data using Euclidean distances, outputs (pseudo- F -statistics) are analogous to traditional least-squares ANOVA, without the same severity of assumptions regarding data distributions and homogeneity of variance (Anderson, 2001; McArdle & Anderson, 2001).

F_v/F_m values for consecutive stress assays were analysed using repeated measures analysis of variance (RM ANOVA) in SPSS v.22 (IBM). The models had three factors: species (2 levels: *L. digitata*, *L. ochroleuca*), temperature (heat emersion and immersion—4 levels: 15°C, 24°C, 28°C & 32°C; cold emersion—2 levels: 15°C & -5°C and; cold immersion—3 levels: 15°C, 4°C & 0°C) and time (11 time points). Relative water content values for heat emersion temperatures were also measured using RM ANOVA with three factors: species (2 levels: *L. digitata* & *L. ochroleuca*), temperature (4 levels: 15°C, 24°C, 28°C, 32°C) and time (9 time points). Where the assumption of sphericity was violated (Mauchly's test) degrees of freedom were corrected, using Greenhouse–Geisser estimates of Sphericity. Also, where significant differences were found (at $p < .05$) post hoc analysis was employed to facilitate interpretation of the RM ANOVA results.

3 | RESULTS

3.1 | Heat shock response

Upregulation of *Hsp70* to the temperature treatments followed a clear bell shaped response for both species (Figure 1). There was no significant interaction between species and temperature (PERMANOVA $F_{(6,56)} = 0.10$ $p = .99$) (Table S1). In both species, thermal set points of the HSR were the same, T_{peak} 24°C and T_{off} 28°C (Figure 1).

3.2 | Consecutive heat shock

3.2.1 | Immersed (heat shock)

Normalised F_v/F_m values varied between time, species and temperature (RM ANOVA: Time \times Species \times Temperature $F_{(10,3,109,6)} = 16.2$, $p < .0001$) (Table S2). A general pattern of a small reduction in F_v/F_m with each 24°C heat shock followed by recovery was observed for both species (Figure 2a,b). In initial heat shock treatments, these were not significantly different from the control. However, by the 3rd heat shock for *L. digitata* and 4th for *L. ochroleuca* values were significantly different to control discs held at 15°C. Whilst not returning to control values after 3 days of recovery both species showed patterns of resilience, indicating that heat stress was reversible.

At 28°C, species showed clear differences in their responses from the first heat shock treatment. Initial treatments (shocks 1–3) only caused minor reductions in F_v/F_m values for *L. ochroleuca* followed by a considerable reduction at shock 4 (Figure 2d). However, there were signs of recovery after each 24 hr period and F_v/F_m levels almost reached those of controls after 3 days recovery. On the other hand, *L. digitata* was only able to recover from its first shock at 28°C. By the fourth cycle values had reached <40% of the control and all discs non-viable 24 hrs after the final shock (Figure 2c).




FIGURE 1 Mean (+1 SE) relative upregulation of *Hsp70* of *Laminaria digitata* and *Laminaria ochroleuca* from St Mawes Bay, UK to 1 hr heat shock treatments ($n = 5$). *Laminaria digitata* normalised to 2 reference genes (ribosomal RNA and Rubisco large subunit), *L. ochroleuca* normalised to 1 reference gene (Rubisco large subunit)

Neither species showed any ability to tolerate heat shocks of 32°C with all discs non-viable by the second cycle (Figure 2e,f).

3.2.2 | Immersed (cold shock)

F_v/F_m values did not vary significantly between species, temperature and time (RM ANOVA: Time \times Species \times Temperature $F_{(6,3,75)} = 1.9$ $p = .07$) (Table S4). At 4°C, there was no pattern of reduced F_v/F_m values pre- or post-treatment (Figure 3a,b). At 0°C, treatments began to affect F_v/F_m values but differences were not significantly different from control values, suggesting that these conditions were non-stressful (Figure 3c,d).

3.2.3 | Emerged (heat shock)

F_v/F_m values varied significantly between species, temperature and time (RM ANOVA: Time \times Species \times Temperature $F_{(8,7,92,3)} = 2.27$ $p = .03$) (Table S3). At 24°C both species showed similar patterns of response with an almost full recovery from exposures 1 and 2. However, after the third exposure recovery was impeded. A pattern of slow recovery was seen in both species yet neither reached close to control values after 72 hrs (Figure 4a,b).

At 28°C a similar pattern of depression of F_v/F_m values followed by recovery was seen in both species for the first three cycles. However, recovery from the fourth cycle was only evident for *L. digitata* (Figure 4c) with values for *L. ochroleuca* still declining after 72 hrs (Figure 4d).

At 32°C a similar pattern to those seen above was observed for the first two cycles, with recovery occurring after 24 hrs. However, after the third cycle recovery was considerably reduced. During the recovery period F_v/F_m values continued to decline in both species (Figure 4e,f). This was most marked in *L. ochroleuca*.




FIGURE 2 Mean (± 1 SE) F_v/F_m values (normalised to 1) for *Laminaria digitata* (left) and *Laminaria ochroleuca* (right) from St Mawes Bay, UK in response to consecutive 1 hr immersion exposures at (a + b) 24°C, (c + d) 28°C and (e + f) 32°C ($n = 5$). Closed symbols are taken immediately after treatments. Initial open symbol represents F_v/F_m prior to any experimental treatment. Subsequent open symbols are taken 24 hrs after treatments to track daily recovery. Dotted line represents mean F_v/F_m for 15°C controls. + represents significant difference from control for that temperature treatment. * represents significant difference between species for that temperature treatment. Different letters represent significant differences from means within each species and treatment




FIGURE 3 Mean (± 1 SE) F_v/F_m values (normalised to 1) for *Laminaria digitata* (left) and *Laminaria ochroleuca* (right) from St Mawes Bay, UK in response to consecutive 1 hr immersion shocks at (a + b) 4°C and (c + d) 0°C ($n = 5$). Closed symbols are taken immediately after treatments. Initial open symbol represents F_v/F_m prior to any experimental treatment. Subsequent open symbols are taken 24 hrs after treatments to track daily recovery. Dotted line represents 15°C controls




FIGURE 4 Mean (± 1 SE) F_v/F_m values (normalised to 1) for *Laminaria digitata* (left) and *Laminaria ochroleuca* (right) from St Mawes Bay, UK in response to consecutive 1 hr emersion shocks at (a + b) 24°C, (c + d) 28 and (e + f) 32°C ($n = 5$). Closed symbols are taken immediately after treatments. Initial open symbol represents F_v/F_m prior to any experimental treatment. Subsequent open symbols are taken 24 hrs after treatments to track daily recovery. Dotted line represents 15°C controls. + represents significant difference from control. * represents significant difference between species. Different letters represent significant differences from means within each species and treatment

3.3 | Water loss

Water loss varied significantly between time, species and temperature (RM ANOVA: Time \times Species \times Temperature $F_{(9,6,102)} = 2.63$ $p = .007$) (Table S6). Kelp subjected to aerial temperatures followed a clear cycle of reduced relative water content (RWC) after each exposure, followed by an increase after 24 hrs (Figure 5). The magnitude of reduction in RWC increased with increasing temperatures while the ability to recover to pre heat shocked water content decreased. A cumulative reduction in RWC with each consecutive treatment was observed across all temperatures for both species. The rate of water loss was consistently greater in *L. ochroleuca*, which also failed to rehydrate fully after 24 hr recovery. In comparison, *L. digitata* was generally almost fully hydrated after the recovery period (Figure 5).

3.3.1 | Emerged (cold shock)

Laminaria digitata and *L. ochroleuca* exhibited very different tolerances to -5°C emersion stress (RM ANOVA: Time \times Species \times Temperature $F_{(1,7,27,1)} = 8.436$ $p = .002$) (Table S5).

Laminaria digitata showed reduced F_v/F_m values with each exposure recovering to near control levels after 24 hrs recovery for the first two cycles. After the third exposure, values were still depressed after 24 hrs and a subsequent exposure resulted in no further reduction. Following 72 hrs, recovery values had returned to almost that of control values indicating that this freezing stress was reversible (Figure 6a). *Laminaria ochroleuca* was much less tolerant to this treatment with large reductions in F_v/F_m values from the initial exposure which dropped further after 24 hrs recovery. After the four exposures, values were down to 15% ($\pm 12.3\%$ SE) of the control and there were no signs of recovery after 72 hrs (Figure 6b).

4 | DISCUSSION

4.1 | Interspecific difference in consecutive low tide tolerance

The two species exhibited varied responses to consecutive immersed heat shocks with the warmer water species, *L. ochroleuca*, better equipped to deal with higher temperature treatments. At




FIGURE 5 Mean (± 1 SE) relative water content of *Laminaria digitata* (left) and *Laminaria ochroleuca* (right) from St Mawes Bay, UK in response to four consecutive 1 hr emersion exposures at (a + b) 24°C, (c + d) 28°C and (e + f) 32°C ($n = 5$). Initial open symbol represents F_v/F_m prior to any experimental treatment. Closed symbols are taken immediately after treatments. Initial open symbol represents F_v/F_m prior to any experimental treatment. Subsequent open symbols are taken 24 hrs after treatments to track daily recovery. + represents significant difference from control. * represents significant difference between species. Different letters represent significant differences from means within each species and treatment




FIGURE 6 Mean (± 1 SE) F_v/F_m values (normalised to 1) for *Laminaria digitata* (left) and *Laminaria ochroleuca* (right) from St Mawes Bay, UK in response to consecutive 1 hr emersion shocks at -5°C . Closed symbols are taken immediately after treatments. Initial open symbol represents F_v/F_m prior to any experimental treatment. Subsequent open symbols are taken 24 hrs after treatments to track daily recovery. Dotted line represents 15°C controls. + represents significant difference from control. * represents significant difference between species. Different letters represent significant differences from means within each species and treatment

24°C, a progressive erosion of resilience was seen in *L. digitata* that was not observed in *L. ochroleuca*. By the final treatment at 28°C, all *L. digitata* were non-viable whereas *L. ochroleuca* were able to

return to control values after 3 days of recovery. Interestingly, a similar lethal response at 28°C was not observed in *L. digitata* during emersed treatments which is likely a product of evaporative cooling

reducing the tissue temperature or desiccation inducing an ametabolic state. Both species differed in their abilities to tolerate stress related to emersion. Most strikingly, *L. ochroleuca* showed no tolerance to even a single freezing exposure. Indeed, anecdotal evidence supports this with reports that over mild winters, *L. ochroleuca* progressively advances up the shore becoming more abundant at higher shore heights, until a severe ground frost results in rapid die off of in the intertidal (G. Boalch, pers. comm.). *Laminaria ochroleuca* was also less equipped to tolerate elevated aerial temperatures, which seemed related to the greater water loss experienced resulting in more severe desiccation. *Laminaria digitata* has been shown to recover from 60% water loss (Dring, 1982), which was also evident in this study. Whilst desiccation tolerance levels are unavailable for *L. ochroleuca*, it was where water loss fell below c. 60% that recovery was not possible.

Climate change in the UK is predicted to result in less summer cloud, increased air temperatures and sunny days, while winter ground frosts will become more infrequent (Jenkins et al., 2009). This may result in *L. ochroleuca*'s maximum tidal height switching from being cold limited in the winter to desiccation limited in the summer. Therefore, the habitat vacated by *L. digitata* may remain unoccupied by *L. ochroleuca* as conditions will still remain outside of its fundamental niche. However, it should be noted that conditions may change asymmetrically. Night temperatures are rising faster than day temperatures (Davy, Esau, Chernokulsky, Outten, & Zilitinkevich, 2017) so advancement up the shore may be possible at least for a period of time. One habitat where *L. ochroleuca* may be able to expand into is tidal pools. Tidal pools in SW England can already reach >28°C in summer (Martins et al., 2007), a temperature that results in considerable stress to *L. digitata* but not *L. ochroleuca*. Currently, only *L. digitata* occupies these habitats suggesting *L. ochroleuca* is being competitively excluded. As temperatures rise and *L. digitata* becomes less competitively dominant or vacates these tidal pools, *L. ochroleuca* will face few barriers to expanding its realised niche to encompass these habitats.

4.2 | Comparing the heat shock response and F_v/F_m (single exposures)

We observed a clear disparity between the HSR and F_v/F_m as metrics for stress. No interspecific differences in thermal set points of the HSR were found but differences were observed in F_v/F_m to similar single stress treatments. For example, *L. digitata* were able to recover from exposures that represent T_{peak} and T_{off} of the HSR, while *L. ochroleuca* showed no deviation away from control values. This is surprising given that T_{peak} and T_{off} are thought to represent the very upper limits of a species tolerance (Barua & Heckathorn, 2004).

Mismatches between chlorophyll fluorescence and the HSR have been observed before (Jueterbock et al., 2014) and may be due to the photosynthetic apparatus having other protective mechanisms in place, allowing for greater thermotolerance compared to the rest of the cell (Downs, Mueller, Phillips, Fauth, & Woodley, 2000). For example, the small HSP, cp-sHSP, is known to play an important role in protecting photosystem II during heat stress, and production levels

are related to thermotolerance in higher plants (Neta-Sharir, Isaacson, Lurie, & Weiss, 2005; Shakeel, Haq, Heckathorn, & Luthe, 2012). Chlorophyll fluorescence may also differ from Hsp quantification as it measures different aspects of heat stress and not just symptoms of protein damage. For example, stress induced uncoupling of enzymes and metabolic pathways can cause the accumulation of reactive oxygen species that can lead to oxidative stress affecting F_v/F_m values (Liu & Pang, 2010). Plants and seaweeds can upregulate genes with anti-oxidative functions during periods of heat stress to protect the photosystems (Collén, Guisile-Marsollier, Léger, & Boyen, 2007) and as such maintain photosynthetic function. Thus, different pressures and mechanisms may affect the two metrics differently and they should not be used interchangeably. Moreover, measuring the HSR alone is not sufficient to fully determine the levels of stress affecting macrophytes as they are likely to be more tolerant than levels of T_{peak} and even T_{off} would suggest.

4.3 | Single vs. consecutive exposures

Intertidal stress studies have overwhelmingly focussed on absolute limits from single treatments with cumulative effects largely being ignored. We identified clear interspecific differences in tolerances to low tide scenarios that were only apparent after consecutive exposures. For example, if a single treatment had been conducted on *L. digitata* at 28°C, we would conclude that such temperature shocks would be non-lethal as full recovery was attained within 24 hrs. Moreover, when comparing this to *L. ochroleuca* we may conclude they exhibit similar tolerances. However, after the final treatment, all *L. digitata* were non-viable whereas *L. ochroleuca* had returned to near control values. Combined with data from the HSR indicating similar tolerances, measuring a single treatment would lead to potentially misleading interpretations.

Environmental Niche Models (ENMs also called distribution, envelope and bioclimatic models) are the most widely used tools to predict the impact of climate change in species distributions. The majority of ENMs determine where a species can exist in the future based on the conditions they exist in now. However, in recent years there has been an attempt to base ENMs on mechanistic cause and effect relationships with environmental variables (Jordà, Marbà, & Duarte, 2012; Martínez, Arenas, Trilla, Viejo, & Carreño, 2015; Sunday et al., 2012). Modellers seek to link critical temperature thresholds (e.g. CT_{max} and CT_{min}) with current and future warming scenarios. Here we show an erosion of resilience leading to mortality at temperatures lower than CT_{max} with consecutive exposures. Therefore, such models may underestimate the effect of future warming on species distributions, with local extinctions potentially occurring at lower temperatures. This is especially true for intertidal and shallow subtidal species that experience discrete consecutive exposures on a daily basis or during a set of exposures that coincide with low spring tides. In such cases, smaller increases in temperatures or discrete "heatwave" events (Hobday et al., 2016), could result in consecutive exposures lower than CT_{max} but great enough to result in considerable stress and subsequent mortality. Therefore, the effect of consecutive exposures on thermal tolerance should be factored into predictive models wherever possible.

5 | CONCLUSION

Our study provides insight into whether a range expander can replace a range contractor at the interface where they coexist within a biogeographic boundary zone. We show that cumulative stress is a key factor in determining interspecific differences in stress physiology. Such knowledge serves to increase our understanding of the processes driving species redistributions, which is comparatively lacking for marine organisms. However, some caution should be taken when generalising results to the wider region. Although the magnitude of difference between species is quite striking and unlikely to arise from site specific local adaptations such factors cannot be discounted, as only a single site and time point were sampled. Moreover, we have only investigated physiological differences between species and species interactions (e.g. facilitation) have not been accounted for. Such interactions may be important in further elucidating species responses to warming.

ACKNOWLEDGEMENTS

N.G.K. was funded by an Institute of Biological, Environmental and Rural Sciences PhD studentship. IBERS receives strategic funding from the Biotechnology and Biological Sciences Research Council. Funding for P.J.M. and N.G.K. was provided through a Marie Curie Career Integration Grant (PCIG10-GA-2011-303685) and the Aberystwyth University Research Fund (1211Z-01). L.S.H. was funded through an Aberystwyth University research fund PhD studentship. D.A.S. is supported by an Independent Research Fellowship award by the Natural Environment Research Council of the UK (NE/K008439/1).

AUTHORS' CONTRIBUTIONS

N.G.K., P.J.M. and D.C.W. conceived the ideas and designed methodology; N.G.K. and L.S.H. collected the data; N.G.K. and P.J.M. analysed the data; all authors contributed to data interpretation; N.G.K. led the writing of the manuscript, with equal input from all authors. All authors contributed critically to the drafts and gave final approval for publication.

DATA ACCESSIBILITY

Data available from Figshare online repository <https://figshare.com/s/c2ad62334bdec04eb976> (King et al., 2017).

ORCID

Nathan G. King  <http://orcid.org/0000-0003-0907-544X>

Dan A. Smale  <http://orcid.org/0000-0003-4157-541X>

REFERENCES

- Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, 26, 32–46.
- Bartsch, I., Vogt, J., Pehlke, C., & Hanelt, D. (2013). Prevailing sea surface temperatures inhibit summer reproduction of the kelp *Laminaria digitata* at Helgoland (North Sea). *Journal of Phycology*, 49, 1061–1073.
- Barua, D., & Heckathorn, S. A. (2004). Acclimation of the temperature set-points of the heat-shock response. *Journal of Thermal Biology*, 29, 185–193.
- Bolton, J., & Anderson, R. (1987). Temperature tolerances of two southern African Ecklonia species (Alariaceae: Laminariales) and of hybrids between them. *Marine Biology*, 96, 293–297.
- Bolton, J., & Lüning, K. (1982). Optimal growth and maximal survival temperatures of Atlantic *Laminaria* species (Phaeophyta) in culture. *Marine Biology*, 66, 89–94.
- Burrows, M. T., Schoeman, D. S., Buckley, L. B., Moore, P., Poloczanska, E. S., Brander, K. M., ... Halpern, B. S. (2011). The pace of shifting climate in marine and terrestrial ecosystems. *Science*, 334, 652–655.
- Butler, W. L. (1978). Energy distribution in the photochemical apparatus of photosynthesis. *Annual Review of Plant Physiology*, 29, 345–378.
- Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B., & Thomas, C. D. (2011). Rapid range shifts of species associated with high levels of climate warming. *Science*, 333, 1024–1026.
- Christidis, N., Stott, P. A., Brown, S., Hegerl, G. C., & Caesar, J. (2005). Detection of changes in temperature extremes during the second half of the 20th century. *Geophysical Research Letters*, 32, L20716.
- Clarke, K. R., & Gorley, R. N. (2006) PRIMER V6: User manual-tutorial. Plymouth, UK: Plymouth Marine Laboratory.
- Collén, J., Guisle-Marsollier, I., Léger, J. J., & Boyen, C. (2007). Response of the transcriptome of the intertidal red seaweed *Chondrus crispus* to controlled and natural stresses. *New Phytologist*, 176, 45–55.
- Connell, S. (2003). The monopolization of understory habitat by subtidal encrusting coralline algae: A test of the combined effects of canopy-mediated light and sedimentation. *Marine Biology*, 142, 1065–1071.
- Coumou, D., & Rahmstorf, S. (2012). A decade of weather extremes. *Nature Climate Change*, 2, 491–496.
- Davy, R., Esau, I., Chernokulsky, A., Outten, S., & Zilitinkevich, S. (2017). Diurnal asymmetry to the observed global warming. *International Journal of Climatology*, 37, 79–93.
- Doney, S. C., Ruckelshaus, M., Emmett Duffy, J., Barry, J. P., Chan, F., English, C. A., ... Knowlton, N. (2012). Climate change impacts on marine ecosystems. *Annual Review of Marine Science*, 4, 11–37.
- Downs, C. A., Mueller, E., Phillips, S., Fauth, J. E., & Woodley, C. M. (2000). A molecular biomarker system for assessing the health of coral (*Montastraea faveolata*) during heat stress. *Marine Biotechnology*, 2, 533–544.
- Dring, M. (1982). Photosynthesis of intertidal brown algae during and after periods of emersion: A renewed search for physiological causes of zonation. *Marine Ecology Progress Series. Oldendorf*, 8, 301–308.
- Eckman, J. E., Duggins, D. O., & Sewell, A. T. (1989). Ecology of under story kelp environments. Effects of kelps on flow and particle transport near the bottom. *Journal of Experimental Marine Biology and Ecology*, 129, 173–187.
- Edwards, M. S., & Estes, J. A. (2006). Catastrophe, recovery and range limitation in NE Pacific kelp forests: A large-scale perspective. *Marine Ecology Progress Series*, 320, 79–87.
- Eggert, A.. (2012) Seaweed responses to temperature. In C. Wiencke & K. Bischof (Eds.), *Seaweed biology* (pp. 47–66). Berlin, Germany: Springer.
- Engelen, A. H., Espirito-Santo, C., Simoes, T., Monteiro, C., Serrão, E. A., Pearson, G. A., & Santos, R. O. (2008). Periodicity of propagule expulsion and settlement in the competing native and invasive brown seaweeds, *Cystoseira humilis* and *Sargassum muticum* (Phaeophyta). *European Journal of Phycology*, 43, 275–282.
- Evans, R. G. (1948). The lethal temperatures of some common British littoral molluscs. *The Journal of Animal Ecology*, 17, 165–173.
- Filbee-Dexter, K., Feehan, C. J., & Scheibling, R. E. (2016). Large-scale degradation of a kelp ecosystem in an ocean warming hotspot. *Marine Ecology Progress Series*, 543, 141–152.
- Forbes, E. (1853) *A history of british mollusca and their shells: By Edw. Forbes and Sylvan. Hanley, UK: John van voorst.*

- Frydman, J. (2001). Folding of newly translated proteins in vivo: The role of molecular chaperones. *Annual Review of Biochemistry*, 70, 603–647.
- García-Mendoza, E., & Colombo-Pallotta, M. F. (2007). The giant kelp *Macrocystis pyrifera* presents a different nonphotochemical quenching control than higher plants. *New Phytologist*, 173, 526–536.
- Gaston, K. J. (2003). *The structure and dynamics of geographic ranges*. Oxford, UK: Oxford University Press.
- Hampe, A., & Petit, R. J. (2005). Conserving biodiversity under climate change: The rear edge matters. *Ecology Letters*, 8, 461–467.
- Hargrave, M. S., Foggo, A., Pessarrodona, A., & Smale, D. A. (2016). The effects of warming on the ecophysiology of two co-existing kelp species with contrasting distributions. *Oecologia*, 183, 531–543.
- Harley, C. D., Anderson, K. M., Demes, K. W., Jorve, J. P., Kordas, R. L., Coyle, T. A., & Graham, M. H. (2012). Effects of climate change on global seaweed communities. *Journal of Phycology*, 48, 1064–1078.
- Harley, C. D., & Paine, R. T. (2009). Contingencies and compounded rare perturbations dictate sudden distributional shifts during periods of gradual climate change. *Proceedings of the National Academy of Sciences*, 106, 11172–11176.
- Hartl, F. U., & Hayer-Hartl, M. (2002). Molecular chaperones in the cytosol: From nascent chain to folded protein. *Science*, 295, 1852–1858.
- Hartmann, D., Tank, A., & Rusticucci, M. (2013) IPCC fifth assessment report, climate change 2013: The physical science basis. IPCC AR5, 31–39.
- Hawkins, S., Sugden, H., Mieszkowska, N., Moore, P., Poloczanska, E., Leaper, R., ... Thompson, R. (2009). Consequences of climate-driven biodiversity changes for ecosystem functioning of North European rocky shores. *Marine Ecology Progress Series*, 396, 245–259.
- Helmuth, B., Choi, F., Matzelle, A., Torossian, J. L., Morello, S. L., Mislán, K., ... Gilman, S. E. (2016). Long-term, high frequency in situ measurements of intertidal mussel bed temperatures using biomimetic sensors. *Scientific Data*, 3, 160087.
- Hendrick, J. P., & Hartl, F.-U. (1993). Molecular chaperone functions of heat-shock proteins. *Annual Review of Biochemistry*, 62, 349–384.
- Henkel, S. K., & Hofmann, G. E. (2008). Differing patterns of *hsp70* gene expression in invasive and native kelp species: Evidence for acclimation-induced variation. *Journal of Applied Phycology*, 20, 915–924.
- Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C., ... Feng, M. (2016). A hierarchical approach to defining marine heatwaves. *Progress in Oceanography*, 141, 227–238.
- Hutchinson, G. E. (1957). Concluding remarks. *Cold Spring Harbor Symposium on Quantitative Biology*, 22, 415–427.
- Hutchinson, G. (1978). *An introduction to population biology*. New Haven, CT: Yale University Press.
- Jenkins, G., Murphy, J., Sexton, D., Lowe, J., Jones, P., & Kilsby, C. G. (2009). *UK climate projections: Briefing report*. Exeter, UK: Met Office Hadley Centre.
- Jordà, G., Marbà, N., & Duarte, C. M. (2012). Mediterranean seagrass vulnerable to regional climate warming. *Nature Climate Change*, 2, 821–824.
- Jueterbock, A., Kollias, S., Smolina, I., Fernandes, J. M., Coyer, J. A., Olsen, J. L., & Hoarau, G. (2014). Thermal stress resistance of the brown alga *Fucus serratus* along the North-Atlantic coast: Acclimatization potential to climate change. *Marine Genomics*, 13, 27–36.
- King, N. G., Wilcockson, D. C., Webster, R., Smale, D. A., Hoelters, L. S., & Moore, P. J. (2017). Cumulative stress restricts niche filling potential of habitat-forming kelps in a future climate. figshare, <https://figshare.com/s/c2ad62334bdec04eb976>
- Lima, F. P., & Wetthey, D. S. (2012). Three decades of high-resolution coastal sea surface temperatures reveal more than warming. *Nature Communications*, 3, 704.
- Liu, F., & Pang, S. J. (2010). Performances of growth, photochemical efficiency, and stress tolerance of young sporophytes from seven populations of *Saccharina japonica* (Phaeophyta) under short-term heat stress. *Journal of Applied Phycology*, 22, 221–229.
- Lourenço, C. R., Zardi, G. I., McQuaid, C. D., Serrao, E. A., Pearson, G. A., Jacinto, R., & Nicastro, K. R. (2016). Upwelling areas as climate change refugia for the distribution and genetic diversity of a marine macroalga. *Journal of Biogeography*, 43, 1595–1607.
- Lüning, K. (1984). Temperature tolerance and biogeography of seaweeds: The marine algal flora of Helgoland (North Sea) as an example. *Helgoländer Meeresunters*, 38, 305–317.
- Lüning, K., & Freshwater, W. (1988). Temperature tolerance of northeast Pacific marine algae. *Journal of Phycology*, 24, 310–315.
- Lüning, K., & tom Dieck, I. (1990) The distribution and evolution of the Laminariales: North Pacific–Atlantic relationships. In D. J. Garbary & R. South (Eds.), *Evolutionary biogeography of the marine algae of the North Atlantic*, pp. 187–204. Berlin, Germany: Springer-Verlag.
- Martínez, B., Arenas, F., Rubal, M., Burgués, S., Esteban, R., García-Plazaola, I., ... Sousa-Pinto, I. (2012). Physical factors driving intertidal macroalgae distribution: Physiological stress of a dominant furoid at its southern limit. *Oecologia*, 170, 341–353.
- Martínez, B., Arenas, F., Trilla, A., Viejo, R. M., & Carreño, F. (2015). Combining physiological threshold knowledge to species distribution models is key to improving forecasts of the future niche for macroalgae. *Global Change Biology*, 21, 1422–1433.
- Martins, G., Hawkins, S., Thompson, R., & Jenkins, S. (2007). Community structure and functioning in intertidal rock pools: Effects of pool size and shore height at different successional stages. *Marine Ecology Progress Series*, 329, 43–55.
- McArdle, B. H., & Anderson, M. J. (2001). Fitting multivariate models to community data: A comment on distance-based redundancy analysis. *Ecology*, 82, 290–297.
- Meehl, G. A., & Tebaldi, C. (2004). More intense, more frequent, and longer lasting heat waves in the 21st century. *Science*, 305, 994–997.
- Neta-Sharir, I., Isaacson, T., Lurie, S., & Weiss, D. (2005). Dual role for tomato heat shock protein 21: Protecting photosystem II from oxidative stress and promoting color changes during fruit maturation. *The Plant Cell*, 17, 1829–1838.
- Orfanidis, S. (1991). Temperature responses and distribution of macroalgae belonging to the warm-temperate Mediterranean-Atlantic distribution group. *Botanica Marina*, 34, 541–552.
- Parsell, D., & Lindquist, S. (1993). The function of heat-shock proteins in stress tolerance: Degradation and reactivation of damaged proteins. *Annual Review of Genetics*, 27, 437–496.
- Pearson, G. A., Lago-Leston, A., & Mota, C. (2009). Frayed at the edges: Selective pressure and adaptive response to abiotic stressors are mismatched in low diversity edge populations. *Journal of Ecology*, 97, 450–462.
- Pearson, G., Lago-Leston, A., Valente, M., & Serrão, E. (2006). Simple and rapid RNA extraction from freeze-dried tissue of brown algae and seagrasses. *European Journal of Phycology*, 41, 97–104.
- Pereira, T. R., Engelen, A. H., Pearson, G. A., Valero, M., & Serrão, E. A. (2015). Response of kelps from different latitudes to consecutive heat shock. *Journal of Experimental Marine Biology and Ecology*, 463, 57–62.
- Poloczanska, E. S., Brown, C. J., Sydeman, W. J., Kiessling, W., Schoeman, D. S., Moore, P. J., ... Burrows, M. T. (2013). Global imprint of climate change on marine life. *Nature Climate Change*, 3, 919–925.
- Raybaud, V., Beaugrand, G., Goberville, E., Delebecq, G., Destombe, C., Valero, M., ... Gevaert, F. (2013). Decline in kelp in west Europe and climate. *PLoS ONE*, 8, e66044.
- Rosman, J. H., Koseff, J. R., Monismith, S. G., & Grover, J. (2007). A field investigation into the effects of a kelp forest (*Macrocystis pyrifera*) on coastal hydrodynamics and transport. *Journal of Geophysical Research: Oceans*, 112, C02016.
- Rothäusler, E., Gómez, I., Karsten, U., Tala, F., & Thiel, M. (2011). Physiological acclimation of floating *Macrocystis pyrifera* to temperature and irradiance ensures long-term persistence at the sea surface at mid-latitudes. *Journal of Experimental Marine Biology and Ecology*, 405, 33–41.

- Saada, G., Nicastro, K. R., Jacinto, R., McQuaid, C. D., Serrao, E. A., Pearson, G. A., & Zardi, G. I. (2016). Taking the heat: Distinct vulnerability to thermal stress of central and threatened peripheral lineages of a marine macroalga. *Diversity and Distributions*, 22, 1060–1068.
- Schultze, K., Janke, K., Krüß, A., & Weidemann, W. (1990). The macrofauna and macroflora associated with *Laminaria digitata* and *L. hyperborea* at the island of Helgoland (German Bight, North Sea). *Helgoländer Meeresuntersuchungen*, 44, 39–51.
- Seneviratne, S. I., Nicholls, N., Easterling, D., Goodess, C. M., Kanae, S., Kossin, J., ... Rahimi, M. (2012a). Changes in climate extremes and their impacts on the natural physical environment. In C. B. Field, V. Barros, T. F. Stocker, D. Qin, D. J. Dokken, K. L. Ebi, M. D. Mastrandrea, K. J. Mach, G.-K. Plattner, S. K. Allen, M. Tignor, & P. M. Midgley (Eds.), *Managing the risks of extreme events and disasters to advance climate change adaptation* (pp. 109–230). Cambridge, UK and New York, NY: Cambridge University Press.
- Shakeel, S. N., Haq, N. U., Heckathorn, S., & Luthe, D. (2012). Analysis of gene sequences indicates that quantity not quality of chloroplast small HSPs improves thermotolerance in C4 and CAM plants. *Plant Cell Reports*, 31, 1943–1957.
- Simonson, E., Metaxas, A., & Scheibling, R. (2015). Kelp in hot water: II. Effects of warming seawater temperature on kelp quality as a food source and settlement substrate. *Marine Ecology Progress Series*, 537, 105–119.
- Smale, D. A., & Wernberg, T. (2013). Extreme climatic event drives range contraction of a habitat-forming species. *Proceedings Royal Society B*, 280, 20122829.
- Smale, D. A., Wernberg, T., Yunnice, A. L., & Vance, T. (2015). The rise of *Laminaria ochroleuca* in the Western English Channel (UK) and comparisons with its competitor and assemblage dominant *Laminaria hyperborea*. *Marine Ecology*, 36, 1033–1044.
- Smolina, I., Kollias, S., Jueterbock, A., Coyer, J. A., & Hoarau, G. (2016). Variation in thermal stress response in two populations of the brown seaweed, *Fucus distichus*, from the Arctic and subarctic intertidal. *Open Science*, 3, 150429.
- Steneck, R. S., Graham, M. H., Bourque, B. J., Corbett, D., Erlandson, J. M., Estes, J. A., & Tegner, M. J. (2002). Kelp forest ecosystems: Biodiversity, stability, resilience and future. *Environmental Conservation*, 29, 436–459.
- Sunday, J. M., Bates, A. E., & Dulvy, N. K. (2012). Thermal tolerance and the global redistribution of animals. *Nature Climate Change*, 2, 686–690.
- Tom Dieck, I. (1992). North Pacific and North Atlantic digitate *Laminaria* species (Phaeophyta): Hybridization experiments and temperature responses. *Phycologia*, 31, 147–163.
- Tomanek, L. (2010). Variation in the heat shock response and its implication for predicting the effect of global climate change on species' biogeographical distribution ranges and metabolic costs. *Journal of Experimental Biology*, 213, 971–979.
- Tomanek, L., & Somero, G. N. (1999). Evolutionary and acclimation-induced variation in the heat-shock responses of congeneric marine snails (genus *Tegula*) from different thermal habitats: Implications for limits of thermotolerance and biogeography. *Journal of Experimental Biology*, 202, 2925–2936.
- Tomanek, L., & Somero, G. N. (2002). Interspecific and acclimation-induced variation in levels of heat-shock proteins 70 (*hsp70*) and 90 (*hsp90*) and heat-shock transcription factor-1 (HSF1) in congeneric marine snails (genus *Tegula*): Implications for regulation of hsp gene expression. *Journal of Experimental Biology*, 205, 677–685.
- Tuya, F., Cacabelos, E., Duarte, P., Jacinto, D., Castro, J. J., Silva, T., ... Coca, J. (2012). Patterns of landscape and assemblage structure along a latitudinal gradient in ocean climate. *Marine Ecology Progress Series*, 466, 9–19.
- Ugarte, R., Critchley, A., Serdynska, A., & Deveau, J. (2009). Changes in composition of rockweed (*Ascophyllum nodosum*) beds due to possible recent increase in sea temperature in Eastern Canada. *Journal of Applied Phycology*, 21, 591–598.
- Voerman, S. E., Llera, E., & Rico, J. M. (2013). Climate driven changes in subtidal kelp forest communities in NW Spain. *Marine Environmental Research*, 90, 119–127.
- Wernberg, T., Bennett, S., Babcock, R. C., de Bettignies, T., Cure, K., Depczynski, M., ... Hovey, R. K. (2016). Climate-driven regime shift of a temperate marine ecosystem. *Science*, 353, 169–172.
- Wernberg, T., Kendrick, G. A., & Toohey, B. D. (2005). Modification of the physical environment by an *Ecklonia radiata* (Laminariales) canopy and implications for associated foliose algae. *Aquatic Ecology*, 39, 419–430.
- Wernberg, T., Smale, D. A., Tuya, F., Thomsen, M. S., Langlois, T. J., De Bettignies, T., ... Rousseaux, C. S. (2013). An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nature Climate Change*, 3, 78–82.

SUPPORTING INFORMATION

Additional Supporting Information may be found online in the supporting information tab for this article.

How to cite this article: King NG, Wilcockson DC, Webster R, Smale DA, Hoelters LS, Moore PJ. Cumulative stress restricts niche filling potential of habitat-forming kelps in a future climate. *Funct Ecol*. 2018;32:288–299. <https://doi.org/10.1111/1365-2435.12977>