



LJMU Research Online

Maughan, RJ, Watson, P, Cordery, PAA, Walsh, NP, Oliver, SJ, Dolci, A, Rodriguez-Sanchez, N and Galloway, SDR

Sucrose and Sodium but not Caffeine Content Influence the Retention of Beverages in Humans Under Euhydrated Conditions

<http://researchonline.ljmu.ac.uk/id/eprint/16301/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

Maughan, RJ, Watson, P, Cordery, PAA, Walsh, NP, Oliver, SJ, Dolci, A, Rodriguez-Sanchez, N and Galloway, SDR (2018) Sucrose and Sodium but not Caffeine Content Influence the Retention of Beverages in Humans Under Euhydrated Conditions. International Journal of Sport Nutrition &

LJMU has developed [LJMU Research Online](#) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>

1 **Title:** Sucrose and sodium but not caffeine content influence the retention of beverages in
2 humans under euhydrated conditions

3

4 **Abstract**

5 This study systematically examined the influence of carbohydrate (sucrose), sodium and
6 caffeine on the fluid retention potential of beverages under euhydrated conditions, using the
7 beverage hydration index (BHI) method. Three cohorts, each of 12 young, healthy, active men,
8 ingested 1L of beverages containing four different concentrations of a single component
9 (sucrose, sodium or caffeine) in a double blind, crossover manner. Urine output was collected
10 for the subsequent 4-h. Cumulative urine output was lower and net fluid balance were higher
11 after 10% and 20% sucrose beverages than 0% and 5% sucrose beverages ($P<0.05$), and after
12 27mmol/L and 52mmol/L sodium beverages than 7mmol/L and 15mmol/L sodium beverages
13 ($P<0.05$). No difference in urine output or net fluid balance was apparent following ingestion of
14 caffeine at concentrations of 0 - 400 mg/l ($P=0.83$). Consequently, the calculated BHI was
15 greater in beverages with higher sucrose or sodium content, but caffeine had no effect. No
16 difference was observed in arginine vasopressin or aldosterone between any trials. These data
17 highlight that the key drivers promoting differences in the fluid retention potential of
18 beverages when euhydrated are energy density, likely through slowed fluid delivery to the
19 circulation (carbohydrate content effect), or electrolyte content through improved fluid
20 retention (sodium content effect). These data demonstrate that beverage carbohydrate and
21 sodium content influence fluid delivery and retention in the 4-h after ingestion, but caffeine up
22 to 400mg/L does not. Athletes and others can use this information to guide their daily
23 hydration practices.

24

25 **Keywords:** carbohydrate, diuresis, electrolytes, gastric emptying

26 **Introduction**

27 Several factors are known to affect maintenance or restoration of fluid balance. The volume
28 and composition of ingested fluids are obviously key in meeting daily water needs and in
29 restoration of fluid balance following exercise (Shirreffs & Maughan, 2000). Although the
30 impact of beverage composition on rehydration has been studied widely over the past 25
31 years, it has been focused around restoration of fluid balance following exercise heat stress-
32 induced dehydration. Responses to fluid intake under euhydrated rested conditions have not
33 been widely explored, though a Beverage Hydration Index (BHI) has recently been proposed to
34 summarise such effects (Maughan et al., 2016) and recently it was demonstrated that body
35 mass and sex do not influence the BHI (Sollanek et al., 2018).

36 Under resting euhydrated conditions, it appears that the carbohydrate, protein, and
37 electrolyte content of ingested beverages are key to influencing subsequent urine production,
38 and thus fluid retention (Maughan et al., 2016). Ingested fluids with a high-energy content
39 (such as milk and fruit juice), as well as those with high electrolyte content (such as milk, fruit
40 juice, and oral rehydration solution(ORS)) promote longer-term retention of the ingested
41 volume (Maughan et al., 2016). These differences in fluid retention are likely due to
42 mechanisms involving both fluid delivery to the circulation (Calbet & Holst, 2004; Mahe et al.,
43 1992) and effect of electrolytes (particularly sodium) on expansion of blood volume and
44 plasma osmolality (Heer et al., 2000). Energy content and osmolality of beverages are known
45 to influence the rate of gastric emptying (Hunt & Stubbs, 1975; Vist & Maughan, 1994, 1995).

46 In addition, glucose and electrolyte composition and osmolality affect intestinal water
47 transport (Schedl et al., 1994; Gisolfi et al., 1992; Shi et al., 1995). Furthermore, the
48 electrolyte content of drinks also affects the retention of fluid within the extracellular or
49 intracellular fluid compartments (Leiper, 2015). Diuretic agents, such as caffeine and alcohol,
50 have little influence on hydration status and fluid loss/retention if taken in small quantities

51 (Armstrong et al., 2005; Maughan et al., 2016; Roti et al., 2006; Seal et al., 2017; Shirreffs &
52 Maughan, 1997). These outcomes have potentially important implications for guidance to
53 individuals/athletes around the ability to retain fluids for longer; particularly during periods
54 when there may be limited access to beverages and when access to facilities for urination are
55 restricted, e.g. when travelling.

56

57 To date, there have been no systematic evaluations of the effect of key beverage components
58 on the retention of beverages during rested euhydrated conditions. For example, the dose of
59 caffeine administered is likely to be key, as doses of caffeine up to 452mg may not induce a
60 significant diuresis vs. matched volumes of water in habitual caffeine users (Armstrong et al.,
61 2005; Killer et al., 2014; Maughan & Griffin, 2003). Recent evidence suggests that only high
62 doses >500mg of caffeine may induce diuresis (Seal et al, 2017) but no systematic evaluation
63 of caffeine dose on fluid balance has been conducted under standardized euhydrated
64 conditions. Furthermore, one study has examined the influence of carbohydrate content of
65 drinks (3% vs 6% carbohydrate) on fluid delivery / retention at rest without prior exercise in
66 mildly dehydrated participants. Over a short follow-up period of only 1-h, no differences were
67 noted for proportion of fluid volume retained between trials (Logan-Sprenger & Spriet, 2013).

68 A recent investigation examined the hydration potential of an amino acid based ORS, a glucose
69 containing ORS and a sports drink and it was demonstrated that the electrolyte content is the
70 primary driver of the fluid retention potential of beverages (Sollanek et al., 2018) . These
71 studies provide some insight but did not systematically examine dose-response effects of
72 different beverage components.

73

74 Thus, to date there has been no systematic assessment of key components, such as
75 carbohydrate, caffeine, and sodium content, on the ability to retain fluid of beverages under
76 euhydrated conditions.

77

78 Therefore, the objective of the present study was to explore the dose-response effects of
79 individual beverage components (sodium, sucrose and caffeine) on the hydration potential of
80 beverages, expressed as the BHI, when ingested under standardized euhydrated conditions. By
81 characterizing the effects of these individual components, we aimed to provide further insight
82 into the factors that determine the BHI response. We hypothesized that increasing the
83 content of sodium and sucrose would increase the ability to retain fluid of beverages
84 expressed as the BHI, while graded caffeine doses within the range commonly ingested (up to
85 400 mg) would have little effect.

86

87 **Methods**

88 *General Study Design*

89 Three laboratories (Loughborough, Bangor and Stirling Universities) collaborated to complete
90 this study. At each site, 12 healthy, weight-stable, active men aged 18-35 years were recruited
91 (n=36 total, **Table 1, Figure 1A**). Participants with a history of cardiovascular, renal,
92 musculoskeletal, or metabolic diseases, as determined from a pre-participation health screen
93 questionnaire, were excluded. Using the experimental approach reported previously
94 (Maughan et al., 2016), each site compared the effect of a control beverage and beverages
95 containing three levels of a single component on post-ingestion fluid balance; Loughborough-
96 caffeine, Stirling-sucrose, Bangor-sodium. Briefly, all urine passed over the 4-h post-ingestion
97 period was collected and expressed as a fraction of that on the water trial. Participants
98 recorded their diet including fluid intake (household measures technique; (Marr, 1971)) and

99 any exercise performed in a diary, over the 2-days before the first trial and referred to this
100 diary to replicate this diet/fluid intake and exercise before the three subsequent visits.
101 Participants were asked not to perform any strenuous exercise or consume alcoholic
102 beverages in the 24-h preceding trials. Compliance was verified verbally with the participants
103 on arrival at the laboratory. Approval for the study was obtained from each of the local Ethics
104 Committees, in accordance with the Declaration of Helsinki (2013). All participants provided
105 written informed consent before participation.

106

107 *Experimental Procedures*

108 Following an overnight fast of ≥ 8 -h, participants emptied their bladder upon waking and
109 retained an aliquot. One hour before arriving at the laboratory, volunteers ingested 500ml of
110 still water (Highland Spring™, Perthshire, UK) over the course of 15min. Upon arrival in the
111 laboratory, volunteers remained seated for 20min. A 20G 1.25" cannula (Becton Dickinson
112 Infusion Therapy Systems Inc., USA) was introduced into an antecubital vein and a blood
113 sample was collected. Participants were then asked to void their bladder and bowels before
114 measurement of body mass (underwear only) to the nearest 50g. Participants then steadily
115 ingested 1L divided in 2 aliquots (every 15min) of the assigned test beverage over a period of
116 30min. At the end of the 30min drinking period, a blood sample was drawn and participants
117 emptied their bladder. This procedure was repeated at hourly intervals, until 4-h post-
118 ingestion. Volunteers remained seated during the drinking period and during the post-
119 ingestion period. Participants stood up when they were asked to empty their bladder or if they
120 needed to void before the collection time point. After the final urine sample was collected,
121 near-nude body mass was recorded again. (**Figure 1B**)

122

123 *Beverages*

124 The control beverage at all sites consisted of still water (Highland Spring™, Perthshire, UK) with
125 added sugar-free fruit-flavoured concentrate (Tesco Stores, UK). This same beverage, with the
126 addition of three levels of a single beverage component, was administered in a randomized,
127 counter-balanced and double-blind manner; Loughborough 50, 200 and 400mg per L of
128 caffeine (BDH, Leicestershire, UK), Stirling 50, 100 and 200g per L of sucrose (British Sugar Ltd,
129 UK), Bangor 15, 27 and 52mmol/L of Na, as sodium chloride (Glacia Fine 60, British Salt Ltd,
130 UK). The control beverage contained 7mmol/L Na and 0.8 g/L of sugar (due to the addition of
131 fruit squash) and was chosen instead of plain water to blind participants to the control trial.
132 The osmolalities of the four beverages administered at Loughborough were 44 (control, 0mg
133 caffeine/L), 43 (50mg caffeine/L), 44 (200mg caffeine/L) and 44mOsmol/kg (400mg caffeine/L),
134 at Stirling were 46 (control, 0.8g/L sucrose), 205 (50g/L sucrose), 386 (100g/L sucrose) and
135 808mOsmol/kg (200g/L sucrose); and at Bangor were 33 (control, 7mmol/L Na), 54 (15mmol/L
136 Na), 85 (27mmol/L Na) and 138mOsmol/kg (52mmol/L Na). Test beverages were stored at a
137 standard refrigerated temperature (4-6 °C) until serving.

138

139 *Urine and blood collection, storage and analysis*

140 Collection, handling, and storage of urine and blood samples were undertaken in accordance
141 with the Human Tissues Act. Stored samples were discarded once analysis was completed.

142

143 All urine collected during the study was passed into a 1L plastic container. The volume of each
144 urine pass was determined by measuring the mass on an electronic balance, assuming a
145 specific gravity of 1.00. From each urine pass, a 5ml aliquot was collected and stored at 4°C.

146 Urine osmolality was measured using freezing-point depression method (Gonotec Osmomat,
147 Germany at Loughborough and Bangor and Roehbling, Camlab, UK at Stirling) within 48-h of
148 collection.

149

150 11mL blood samples were drawn into dry syringes and immediately dispensed into a 5mL
151 serum tube, and 1mL and 5mL EDTA tubes. At Stirling, duplicate 100 µL aliquots of whole
152 blood were rapidly deproteinised in Eppendorf tubes containing 1 mL of ice-cold 0.3 N
153 perchloric acid. These samples were centrifuged and the resulting supernatant used to
154 determine blood glucose concentrations (Glucose oxidase method, Instrumentation
155 Laboratory, Italy).

156

157 Whole blood in the serum tube was allowed to stand for 1-h at room temperature to clot
158 before centrifugation (10min, 4°C, 2000-3000g). Serum was dispensed and stored at 4°C for
159 measurement of osmolality by freezing-point depression and sodium by flame-photometry
160 (Bangor). A further serum aliquot was stored at - 80°C for measurements of aldosterone and
161 arginine vasopressin concentrations by enzyme-linked immunosorbent assay (Enzo Life
162 Sciences, Lausen, Switzerland) and caffeine concentrations by HPLC (Loughborough; Holland et
163 al., 1998)).

164

165 *Beverage hydration index (BHI) calculation*

166 The beverage hydration index (BHI) (Maughan, et al., 2016) was obtained by dividing the total
167 urine output over a period of time for the control beverage by the total urine output for the
168 same period of time after the test beverage was ingested.

$$169 \quad BHI = \frac{\text{Total urine output when control beverage ingested (L)}}{\text{Total urine output when test beverage ingested (L)}}$$

170

171

172 *Data and statistical analysis*

173 Participant characteristics at each institution were compared by one-way ANOVA. Pre-drink
174 hydration status, as assessed by body mass, serum and urine osmolality, was compared by
175 repeated-measures ANOVA. For each beverage component studied the cumulative urine mass,
176 net fluid balance and blood parameters were compared each hour and between different
177 beverage doses by 2-way repeated-measures ANOVA. Significant main effects and interactions
178 were further explored by Tukey's multiple-comparison tests. BHI values were not normally
179 distributed and therefore statistical comparison between beverages was made by Friedman
180 test with significant effects further explored by Dunn's multiple comparison tests. The
181 meaningfulness of differences observed was calculated using 95% CI of differences between
182 means and Cohen's d effect size (Cohen, 1988). All statistical analyses were completed with
183 the use of a statistical software package (GraphPad Prism version 6 for Windows). Statistical
184 significance was accepted at $P < 0.05$.

185

186 Sample size was based on a minimally important difference using 80% power and a two-tailed
187 alpha level of 0.05. Hypothesized effect size was 0.81, calculated from the difference between
188 estimated mean cumulative urine output (minimally important difference of 168mL)
189 (Maughan, et al., 2016) with a pooled SD of 206ml giving an estimated sample size required of
190 $n=12$ per site.

191

192 **Results**

193 Forty participants were recruited: loss to follow-up occurred because of vomiting after
194 beverage ingestion ($n=2$), or because of voluntary withdrawal from the study ($n=2$), resulting in
195 $n=36$ participants, 12 at each site.

196

197 *Pre-drink ingestion hydration status*

198 On each trial, pre-ingestion hydration status indicated euhydration (**Table 2**). The coefficient of
199 variation (CV) for initial body mass was 0.6%, 0.8% and 0.6% for all sucrose, sodium and
200 caffeine trials, respectively. The CV for initial serum osmolality was 0.7%, 1.0% and 0.7% for all
201 sucrose, sodium and caffeine trials, respectively. The CV for initial urine osmolality was 37%,
202 39% and 24% for all sucrose, sodium and caffeine trials, respectively.

203

204 *Blood glucose, serum sodium and plasma caffeine responses*

205 Blood glucose concentration was greater after ingesting beverages containing sucrose (**Figure**
206 **2A**, $P < 0.01$). Up to 1-h after beverage ingestion, blood glucose remained higher after the 20%
207 sucrose beverage than the 0% and 5% beverages. Blood glucose was then similar between
208 beverages for the remainder of the 4-h with exception of the 10% sucrose beverage being
209 lower than the 0% and 20% beverages at 2-h. Serum sodium was not changed after ingesting
210 beverages of different sodium contents (**Figure 2B**). Plasma caffeine content increased in a
211 dose-dependent manner (**Figure 2C**, $P < 0.01$).

212

213 *Urine output and fluid balance responses to sucrose*

214 Immediately after ingesting the different sucrose beverages, urine mass was similar ($P = 0.12$).
215 Cumulative urine output was lower and net fluid balance higher at 1-h, 2-h and 3-h after
216 ingestion of the 10% and 20% sucrose beverages than the 0% and 5% sucrose beverages
217 (**Figures 3A & 3B**, $P < 0.05$). Throughout the 4-h period, cumulative urine output was lower and
218 net fluid balance higher after the 20% sucrose beverage than the 0%, 5% and 10% beverage
219 ($P < 0.05$). The effect sizes at 2-h compared with the 0% beverage were 1.46 for the 20%
220 sucrose beverage and 0.73 for the 10% sucrose beverage. The mean differences in urine
221 output compared with the 0% beverage were 500g (95%CI: 399, 601g) for the 20% sucrose
222 beverage and 189g for the 10% sucrose beverage (95%CI: 87, 290g).

223

224 *Urine output and fluid balance responses to sodium*

225 One hour after ingesting different sodium beverages urine mass was similar ($P = 0.30$), but 2-
226 h, 3-h, 4-h after ingestion cumulative urine output was lower and net fluid balance higher after
227 the 27mmol/L and 52mmol/L sodium beverages than the 7mmol/L and 15mmol/L beverages
228 (**Figures 3C & 3D, $P < 0.05$**). The effect sizes at 3-h compared with the 7mmol/L beverage were
229 1.06 for the 52mmol/L beverage and 0.87 for the 27mmol/L beverage. The mean differences
230 compared with the 7mmol/L beverage were 372g (95%CI: 228, 516g) for the 52mmol/L sodium
231 beverage and 300g (95%CI: 156, 444g) for the 27mmol/L sodium beverage. These differences
232 also exceeded the 3-h cumulative urine output and net fluid balance CV.

233

234 *Urine output and fluid balance responses to caffeine*

235 Urine mass and net fluid balance were similar throughout the 4-h period on all trials after the
236 ingestion of drinks with different caffeine content (**Figures 3E&3F, $P = 0.83$**).

237

238 *Beverage Hydration Index*

239 Based on our previous observations, a calculated BHI exceeding twice the CV of the BHI index
240 can be considered as meaningful, representing a better fluid retention (Maughan et al., 2016).
241 BHI was greater in drinks with higher sucrose and sodium content, but was not affected by
242 caffeine content (**Figure 4, $P < 0.05$**). After 1-h, 2-h, 3-h and 4-h, 20% sucrose beverage had
243 higher BHI than control (0% sucrose beverage) and at 2-h and 3-h was higher than 5% sucrose
244 beverage ($P < 0.05$). After 2-h, 3-h and 4-h the 27mmol/L and 52mmol/L sodium beverages had
245 higher BHI than the control trial (**Figure 4A&4B, all differences $P < 0.05$**).

246

247 *Fluid-regulation and redistribution*

248 Throughout the 4-h period, concentrations of aldosterone and arginine vasopressin were
249 similar irrespective of the sucrose, sodium or caffeine content of beverages (**Table 3**).
250 Immediately after and in the first hour after ingestion of 10% and 20% sucrose content
251 beverages, serum osmolality increased, and was different to control and to 5% sucrose
252 beverage ($P<0.05$), while it was relatively unchanged and similar after 0% and 5% sucrose
253 beverage ingestion (**Figure 5A**). In contrast, immediately after ingestion of sodium beverages,
254 serum osmolality decreased but to a less extent of 52mmol/L sodium beverage in comparison
255 with the control (**Figure 5B, $P<0.05$**). Osmolality was not measured in caffeine trials.

256

257 **Discussion**

258 In the present study cumulative urine output was lower and net fluid balance higher 4-h after
259 the ingestion of the 10% and 20% sucrose beverages than after the ingestion of the 0% and 5%
260 sucrose beverages. A similar response was observed with 27 mmol/L and 52 mmol/L sodium
261 beverages compared to the 7 mmol/L and 15 mmol/L beverages. However, no differences in
262 urine mass or net fluid balance were apparent 4-h following the ingestion of different caffeine
263 contents. These observations are consistent with our initial hypotheses and demonstrate that
264 factors affecting fluid delivery (sucrose content) and retention (sodium content) are
265 dependent upon the dose contained within ingested beverages. These data also demonstrate
266 that caffeine up to 400 mg/L has no impact upon hydration potential or the ability to retain
267 fluid of beverages.

268

269 In our previous work (Maughan et al., 2016), we were able to quantify the hydration potential
270 of commercially-available drinks using a beverage hydration index (BHI). The BHI was
271 postulated to be related to energy density and electrolyte composition, both of which can
272 affect fluid delivery and retention. However, combinations of key components (e.g.

273 macronutrients, electrolytes and caffeine) at different doses could influence gastric emptying,
274 intestinal absorption, and fluid retention characteristics. The results of the present study
275 reveal that, in comparison to control beverage, under euhydrated conditions a sucrose content
276 of up to 5%, a caffeine content of up to 400mg/L, and a sodium content of up to 15mmol/L all
277 have no effect on the BHI. However, 10% and 20% sucrose beverages, and beverages
278 containing 27mmol/L and 52mmol/L sodium result in reduced diuresis. Given that these test
279 drinks were examined under euhydrated conditions, the reduced urine output likely occurred
280 due to mechanisms involving a combination of altered gastric emptying (Hunt & Stubbs, 1975)
281 and intestinal absorption (Leiper, 2015). Furthermore, the electrolyte content has potential
282 effects on fluid retention independent of hormonal controls (Schedl & Clifton, 1963).

283

284 *Gastric emptying, intestinal absorption and renal excretion of fluids*

285 Early studies demonstrated that the addition of sodium to test drinks with low glucose content
286 increased the rate of gastric emptying (Hunt & Pathak, 1960) and intestinal absorption (Phillips
287 & Summerskill, 1967). Other studies demonstrated that glucose at >4% solution content
288 reduced the rate of gastric emptying compared to water, that warm/hot fluids reduced gastric
289 emptying compared to cold beverages, and that faster initial emptying rates were reached
290 with higher bolus volumes (Costill & Saltin, 1974; Hunt & Macdonald, 1954; Vist & Maughan,
291 1994, 1995). Applying these observations to the current study it can be proposed that gastric
292 emptying rate would be increased with an increasing sodium content of beverages (above 33
293 mmol/L), reduced with an increasing energy/carbohydrate content (above 4-5%
294 carbohydrate), and likely remain unchanged by increasing caffeine content (up to 269 mg).
295 Indeed, these largely reflect the reported observations in the present study.

296

297 Intestinal perfusion studies reveal that hypertonic solutions (>300mOsm/kg) result in transient
298 net water secretion into the intestinal lumen whereas hypotonic solutions (<260mOsm/kg)
299 stimulate net water absorption (Hunt et al., 1992). High carbohydrate solutions with high
300 osmolality will therefore delay gastric emptying, slow delivery of fluid to the intestine, and
301 cause net water secretion into the intestinal lumen. Water absorption appears to be
302 independent of carbohydrate at concentrations up to 6% (Gisolfi et al., 1992). Applying these
303 observations to the present study would suggest that more concentrated sucrose solutions
304 ($\geq 10\%$) would likely slow gastric emptying result in transient net water secretion into the
305 intestinal lumen. The effect of increasing the sodium content upon the ability to retain fluid of
306 beverages suggests an initial fast gastric emptying inducing increase in intestinal water and
307 sodium transport, and subsequently greater retention of the fluid in the body water pool. The
308 decrease in serum osmolality observed following beverage ingestion supports these
309 assertions.

310

311 The principal determinant of permeability, and consequently of water reabsorption, in the
312 collecting ducts of the kidneys is arginine vasopressin (AVP) (Bourque, 2010). Aldosterone,
313 produced by the adrenal cortex, also stimulates sodium reabsorption in the cortical collecting
314 ducts (Stanhewicz & Kenney, 2015). In the present study, the responses of aldosterone and
315 AVP to fluid ingestion were similar regardless of the content of sucrose, sodium or caffeine
316 within the beverages. AVP and aldosterone also did not change over time during the ingestion
317 or follow-up period. Thus, in the present work it can be concluded that differences in urine
318 output between sucrose beverages and between sodium-containing beverages are not
319 influenced by differences in renal water or sodium excretion. Thus, by studying participants in
320 a euhydrated state we have been able to isolate effects on fluid delivery/retention while
321 removing potential interaction of hormonal controls. The differences in 2-h cumulative urine

322 output and in net fluid balance observed in the sucrose and in the sodium trials can be
323 considered meaningful as they exceeded the CV calculated previously (Maughan et al., 2016)
324 and the minimally important difference of 168mL calculated a priori.

325

326 *Caffeinated beverages and hydration*

327 Caffeine is an adenosine receptor antagonist reducing fractional sodium reabsorption in the
328 proximal tubule and in the distal nephron (Shirley et al., 2002) which could lead to increased
329 renal water loss. Previous studies exploring the effect of administering different doses of
330 caffeine have observed increased urine volume only when participants ingested 360 mg of
331 caffeine (Passmore et al., 1987), 6 mg/kg of caffeine (Seal et al., 2017) or 624 mg (Neuhauser
332 et al., 1997). In the present study, no difference in urine volume was noted following any of
333 the doses of caffeine administered. This suggests that sodium excretion was not influenced by
334 caffeine in our participants. Unfortunately, sodium excretion in urine was not determined in
335 our trials to enable confirmation of this proposal. The lack of effect of all the caffeine doses
336 studied in the present study supports and adds to earlier observations on caffeine dose. Thus,
337 caffeinated beverages (containing up to 400mg of caffeine) can contribute to daily total fluid
338 intake targets without negative effects on fluid balance.

339

340 *Practical Perspectives / Study Limitations*

341 This study provides further evidence that the sodium content of a beverage is likely to be a
342 main driver for improved fluid delivery and retention, while high carbohydrate content likely
343 delays fluid delivery and increases the serum osmolality, and caffeine up to 400mg has no
344 impact on diuresis 4-h after the beverage ingestion. These mechanistic observations can
345 provide useful information for athletes as their teams can develop a fluid intake strategy for
346 when there is limited access to fluid or when the access to facilities to urinate is restricted (e.g.

347 when the athletes are travelling) The outcomes of the present study require further
348 exploration in other groups such as older adults who have a reduced ability to alter renal
349 water excretion. Future studies also should examine the effects of other macro- and micro-
350 nutrients on the hydration potential of beverages.

351

352 **Acknowledgements**

353 The authors acknowledge the contribution of Dr Lewis James towards data collection
354 undertaken at Loughborough University. This study was funded in part by a grant from the
355 European Hydration Institute. The European Hydration Institute had no role in the design,
356 analysis or writing of this article. Prof Maughan was Chair of the Scientific Advisory Board of
357 the European Hydration Institute. Dr Phil Watson has received funding in the past 3 years from
358 the European Hydration Institute for other hydration-related research. No other authors
359 declare a conflict of interest. R.J.M. conceived the project, R.J.M., P.W., P.A.A.C., N.P.W., S.J.O.,
360 N.R.S. and S.D.R.G. developed the overall research plan. P.W., N.P.W. and S.D.R.G. had study
361 oversight. P.A.A.C., AD and N.R.S. conducted the research and analyzed the samples. S.J.O. and
362 N.P.W. performed the statistical analysis. R.J.M., P.W., N.P.W. and S.D.R.G. wrote the paper
363 with P.A.A.C., S.J.O. and N.R.S. S.D.R.G. had primary responsibility for the final content. All the
364 authors approved the final version of the paper.

365

366

References

- Armstrong, L.E., Pumerantz, A.C., Roti, M.W., Judelson, D.A., Watson, G., Dias, J.C., Sökmen, B., Casa, D.J., Maresh, C.M., Lieberman, H. & Kellogg, M. (2005). Fluid, electrolyte, and renal indices of hydration during 11 days of controlled caffeine consumption. *International Journal of Sport Nutrition & Exercise Metabolism* 15(3), 252-265. [PubMed](#)
- Bourque, C.W., (2008). Central mechanisms of osmosensation and systemic osmoregulation. *Nature Reviews Neuroscience* 9, 519-531. [doi:10.1038/nrn2400](https://doi.org/10.1038/nrn2400)
- Calbet, J.A. & Holst, J.J. (2004). Gastric emptying, gastric secretion and enterogastrone response after administration of milk proteins or their peptide hydrolysates in humans. *European Journal of Nutrition* 43(3), 127-139. [PubMed](#) [doi: 10.1007/s00394-004-0448-4](https://doi.org/10.1007/s00394-004-0448-4)
- Cohen, J. (1988). *Statistical power analysis for the behavioral sciences*. 2nd ed. Hillsdale, NJ: Erlbaum. [doi: 10.4324/9780203771587](https://doi.org/10.4324/9780203771587)
- Costill, D.L. & Saltin, B. (1974). Factors limiting gastric emptying during rest and exercise. *Journal of Applied Physiology* 37(5), 679-683. [PubMed](#)
- Gisolfi, C.V., Summers, R.W., Schedl, H.P. & Bleiler, T.L. (1992). Intestinal water absorption from select carbohydrate solutions in humans. *Journal of Applied Physiology* 73(5), 2142-2150. [PubMed](#)
- Heer, M., Baisch, F., Kropp, J., Gerzer, R. & Drummer, C. (2000). High dietary sodium chloride consumption may not induce body fluid retention in humans. *American Journal of Physiology Renal Physiology* 278(4), F585-595. [PubMed](#)
- Holland, D.T., Godfredsen, K.A., Page, T. & Connor, J.D. (1998). Simple high-performance liquid chromatography method for the simultaneous determination of serum caffeine and paraxanthine following rapid sample preparation. *Journal of Chromatography B:*

Biomedical Sciences and Applications 707(1-2), 105-110. [PubMed doi: 10.1016/s0378-4347\(97\)00590-2](#)

Hunt, J.B., Elliott, E.J., Fairclough, P.D., Clark, M.L. & Farthing, M.J. (1992). Water and Solute Absorption from Hypotonic Glucose-Electrolyte Solutions in Human Jejunum. *Gut* 33(4), 479-483. [PubMed doi: 10.1136/gut.33.4.479](#)

Hunt, J.N. & Macdonald, I. (1954). The influence of volume on gastric emptying. *The Journal of Physiology* 126(3), 459-474. [PubMed doi: 10.1113/jphysiol.1954.sp005222](#)

Hunt, J.N. & Pathak, J.D. (1960). The osmotic effects of some simple molecules and ions on gastric emptying. *The Journal of Physiology* 154(2), 254-269. [PubMed doi: 10.1113/jphysiol.1960.sp006577](#)

Hunt, J.N. & Stubbs D.F. (1975). The volume and energy content of meals as determinants of gastric emptying. *The Journal of Physiology* 245(1), 209-225. [PubMed doi: 10.1113/jphysiol.1975.sp010841](#)

Killer, S.C., Blannin, A.K. & Jeukendrup, A.E. (2014). No evidence of dehydration with moderate daily coffee intake: a counterbalanced cross-over study in a free-living population. *PLoS One* 9(1), e84154. [PubMed doi: 10.1371/journal.pone.0084154](#)

Leiper, J.B. (2015). Fate of ingested fluids: factors affecting gastric emptying and intestinal absorption of beverages in humans. *Nutrition Reviews* 73(2), 57-72. [PubMed doi:10.1093/nutrit/nuv032](#)

Logan-Sprenger, H.M. & Spriet, L.L. (2013). The acute effects of fluid intake on urine specific gravity and fluid retention in a mildly dehydrated state. *Journal of Strength and Conditioning Research* 27(4), 1002-1008. [PubMed](#)

Mahe, S., Huneau, J.F., Marteau, P., Thuillier, F. & Tome, D. (1992). Gastroileal nitrogen and electrolyte movements after bovine milk ingestion in humans. *American Journal of Clinical Nutrition* 56(2), 410-416. [PubMed](#)

- Marr, J.W. (1971). Individual dietary surveys: purposes and methods. *World Review of Nutrition and Dietetics* 13, 105-164. [PubMed doi: 10.1159/000391884](#)
- Maughan, R.J. & Griffin, J. (2003). Caffeine ingestion and fluid balance: a review. *Journal of Human Nutrition and Dietetics* 16(6), 411-420. [PubMed doi: 10.1046/j.1365-277x.2003.00477.x](#)
- Maughan, R.J., Watson, P., Cordery, P.A., Walsh, N.P., Oliver, S.J., Dolci, A., Rodriguez-Sanchez, N & Galloway, S.D.R. (2016). A randomized trial to assess the potential of different beverages to affect hydration status: development of a beverage hydration index. *American Journal of Clinical Nutrition* 103(3), 717-723. [PubMed doi: 10.3945/ajcn.115.114769](#)
- Neuhauser, B., Beine, S., Verwied, S.C. & Luhrmann, P.M. (1997). Coffee consumption and total body water homeostasis as measured by fluid balance and bioelectrical impedance analysis. *Annals of Nutrition & Metabolism* 41(1), 29–36. [doi:10.1159/000177975](#)
- Passmore, A.P., Kondowe, G.B. & Johnston, G.D. (1987). Renal and cardiovascular effects of caffeine: a dose–response study. *Clinical Science* 72(6), 749-756. [PubMed doi: org/10.1042/cs0720749](#)
- Phillips, S.F. & Summerskill, W.H. (1967). Water and electrolyte transport during maintenance of isotonicity in human jejunum and ileum. *The Journal of Laboratory and Clinical Medicine* 70(4), 686-698. [PubMed](#)
- Roti, M.W., Casa, D.J., Pumerantz, A.C., Watson, G., Dias, J.C., Ruffin, K. & Armstrong, L.E. (2006). Thermoregulatory responses to exercise in the heat: Chronic caffeine intake has no effect. *Aviation, Space, and Environmental Medicine* 77(2), 124-129. [PubMed](#)
- Seal, A.D., Bardis, C.D., Gavrieli, A., Grigorakis, P., Adams, J.D., Arnaoutis, G., Yannakoulia, M. & Kavouras, S.A. (2017) Coffee with high but not low caffeine content augments fluid and

- electrolyte excretion at rest. *Frontiers in Nutrition* 4(40), 1-6. [PubMed](#) doi: [10.3389/fnut.2017.00040](https://doi.org/10.3389/fnut.2017.00040)
- Schedl, H.P. & Clifton, J.A. (1963). Solute and water absorption by the human small intestine. *Nature* 199, 1264-1267. [PubMed](#)
- Shi, X., Summers, R.W., Schedl, H.P., Flanagan, S.W., Chang, R. & Gisolfi, C.V. (1995). Effects of carbohydrate type and concentration and solution osmolality on water absorption. *Medicine and Science in Sports and Exercise* 27(12), 1607-1615. [PubMed](#)
- Shirley, D.G., Walter, S.J. & Noormohamed, F.H. (2002). Natriuretic effect of caffeine: assessment of segmental sodium reabsorption in humans. *Clinical Science* 103(5), 461-466. [PubMed](#) doi: [10.1042/cs1030461](https://doi.org/10.1042/cs1030461)
- Shirreffs, S.M. & Maughan, R.J. (1997). Restoration of fluid balance after exercise-induced dehydration: effects of alcohol consumption. *Journal of Applied Physiology* 83(4), 1152-1158. doi: [10.1152/jappl.1997.83.4.1152](https://doi.org/10.1152/jappl.1997.83.4.1152)
- Shirreffs, S.M. & Maughan, R.J. (2000). Rehydration and recovery of fluid balance after exercise. *Exercise and Sport Sciences Reviews* 28(1), 27-32. [PubMed](#)
- Sollanek, K.J., Tsurumoto, M.T., Vidyasagar, S., Kenefick, R.W. & Cheuvront, S.N. (2018). Neither body mass nor sex influences beverage hydration index outcomes during randomized trial when comparing 3 commercial beverages. *American Journal of Clinical Nutrition* 107:544-549. [PubMed](#)
- Stanhewicz, A.E., Kenney, W.L. (2015). Determinants of water and sodium intake and output. *Nutrition Reviews* 73(Suppl_2), 73-82. [PubMed](#) doi: [10.1093/nutrit/nuv033](https://doi.org/10.1093/nutrit/nuv033)
- Vist, G.E. & Maughan, R.J. (1994). Gastric-Emptying of Ingested Solutions in Man - Effect of Beverage Glucose-Concentration. *Medicine & Science in Sports & Exercise* 26(10), 1269-1273. [PubMed](#) doi: [10.1249/00005768-199410000-00014](https://doi.org/10.1249/00005768-199410000-00014)

Vist, G.E. & Maughan, R.J. (1995). The effect of osmolality and carbohydrate content on the rate of gastric emptying of liquids in man. *The Journal of Physiology* 486(Pt 2), 523-531.

PubMed doi: 10.1113/jphysiol.1995.sp020831

Table 1. Participant physical characteristics, measured during the pre-screening consultation, estimated daily water, alcohol and caffeine intake from the food diaries at each of the three study sites and for combined data (all sites).

	Stirling - Sucrose (n = 12)	Bangor - Sodium (n = 12)	Loughborough - Caffeine (n = 12)	All (n = 36)	<i>P</i>
Age (y)	26 ± 6	25 ± 4	27 ± 2	26 ± 4	0.53
Height (cm)	181 ± 7	179 ± 7	178 ± 7	179 ± 7	0.67
Mass (kg)	77.6 ± 9.3	78.2 ± 7.8	77.1 ± 8.9	77.6 ± 8.5	0.95
BMI (kg/m ²)	23.9 ± 2.7	24.6 ± 2.2	24.2 ± 1.5	24.2 ± 2.1	0.75
Water intake (L/d)	1.9 ± 0.3	2.2 ± 0.9	1.9 ± 0.5	2.0 ± 0.6	0.42
Caffeine intake (mg/d)	210 ± 142	180 ± 123	206 ± 176	199 ± 145	0.87
Alcohol intake (g/d)	5 ± 6	4 ± 4	3 ± 2	4 ± 4	0.55

Notes: Data are Mean ± Standard Deviation. Water intake represent fluid from beverages only. Alcohol intake includes all forms of alcoholic beverages. BMI = Body Mass Index.

Table 2. Pre-ingestion hydration status at each of the three study sites.

Stirling – Sucrose (n = 12)					
	0%	5%	10%	20%	P
Body mass (kg)	77.5 ± 9.2	77.5 ± 9.4	77.7 ± 9.1	77.5 ± 9.5	0.70
Serum osmolality* (mmol/kg)	295 ± 3	296 ± 2	296 ± 2	295 ± 2	0.77
Urine osmolality (mmol/kg)	524 ± 323	557 ± 209	488 ± 290	664 ± 332	0.38
Bangor – Sodium (n = 12)					
	7 mmol/L	15 mmol/L	27 mmol/L	52 mmol/L	P
Body mass (kg)	78.2 ± 7.8	78.4 ± 8.1	78.5 ± 7.8	78.1 ± 8.2	0.50
Serum osmolality (mmol/kg)	289 ± 3	290 ± 3	291 ± 4	292 ± 4	0.17
Urine osmolality (mmol/kg)†	520 ± 215	544 ± 232	475 ± 201	513 ± 300	0.82
Loughborough – Caffeine (n = 12)					
	0 mg	50 mg	100 mg	400 mg	P
Body mass (kg)	77.3 ± 10.1	77.5 ± 10.1	77.7 ± 10.1	77.3 ± 10.1	0.26
Serum osmolality (mmol/kg)	287 ± 4	289 ± 5	289 ± 6	290 ± 5	0.05
Urine osmolality (mmol/kg)	441 ± 179	486 ± 144	478 ± 163	519 ± 168	0.48

Notes: Data are presented as Mean ± Standard Deviation.

*osmolality assessment of an identical control solution (mean 292 mmol/kg) at each site indicated that the Roehbling osmometer (Stirling) consistently reported a +4 mmol/kg bias compared with the Gonotec osmometer (Loughborough and Bangor). † n = 11 for Bangor urine osmolality analysis.

Table 3. Mean plasma aldosterone and plasma arginine vasopressin (AVP) responses over the 4-h follow-up period following each test drink ingestion, at each study site.

Stirling – Sucrose (n = 12)					
	0%	5%	10%	20%	P
Aldosterone (pg/ml)	103 ± 31	113 ± 27	100 ± 30	106 ± 34	0.47
AVP (pg/ml)	3.5 ± 0.6	3.4 ± 0.6	3.6 ± 0.6	3.7 ± 0.7	0.50
Bangor – Sodium (n = 12)					
	7 mmol/L	15 mmol/L	27 mmol/L	52 mmol/L	P
Aldosterone (pg/ml)	109 ± 41	126 ± 67	150 ± 59	100 ± 62	0.16
AVP (pg/ml)	3.7 ± 0.7	3.6 ± 0.9	3.8 ± 1.2	3.9 ± 0.8	0.79
Loughborough – Caffeine (n = 12)					
	0 mg	50 mg	200 mg	400 mg	P
Aldosterone (pg/ml)	90 ± 73	99 ± 64	72 ± 64	87 ± 108	0.60
AVP (pg/ml)	3.5 ± 1.4	3.5 ± 1.1	2.9 ± 0.9	3.8 ± 0.9	0.22

Note: Data are presented as Mean ± Standard Deviation.

Figure Legends

FIGURE 1. Experimental design of the study (A) and schematic of experimental protocol (B).

CHO = carbohydrate (sucrose), Na = sodium.

FIGURE 2. Blood glucose (A), serum sodium (B) and plasma caffeine responses (C) after the ingestion of 1 L of various sucrose (A), sodium (B) and caffeine (C) content beverages vs. control. n = 12 observation on each beverage. Beverages with different responses are identified by Tukey's multiple comparison test: a, indicates difference to 0% sucrose (control) or 0 mg caffeine (control) beverage, b, indicates difference to 5% or 50 mg caffeine, c, indicates difference to 10% or 200 mg caffeine. Statistical significance was accepted at $P < 0.05$. The vertical error bar in the top left corner represents the overall mean SD during the 4-h collection.

FIGURE 3. Cumulative urine output and net fluid balance after the ingestion of 1 L of various sucrose (A & B), sodium (C & D) and caffeine (E & F) content beverages. n = 12 observation on each beverage. Beverages with different responses are identified by Tukey's multiple comparison test: a, indicates difference to 0% sucrose (control) or 7 mmol/L sodium (control) beverage; b, indicates difference to 5% sucrose or 15 mmol/L sodium beverage; c, indicates difference to 10% sucrose beverage. Downward arrows indicate the first time when statistical differences were detected between beverages. Statistical significance was accepted at $P < 0.05$. The vertical error bar in the top left corner represents the mean SD during the 4-h collection.


FIGURE 4. Beverage hydration index for various sucrose (A), sodium (B) and caffeine (C) content beverages. n = 12 observation on each beverage. Beverages with different responses are identified by Dunn's multiple comparison test: a, indicates difference to 0% sucrose

(control) or 7 mmol/L sodium (control) beverage; b, indicates difference to 5% sucrose beverage; c, indicates difference to 10% sucrose beverage. Statistical significance was accepted at $P < 0.05$. These are median data with the mean IQR during the 4-h collection represented by the vertical error bar in the top left corner. Downward arrows indicate the first time when statistical differences were detected between beverages.

FIGURE 5. Serum osmolality change after the ingestion of 1 L of various sucrose (A) and sodium (B) beverages. $n = 12$ observation on each beverage. Beverages with different responses are identified by Tukey's multiple comparison test: a, indicates difference to 0% sucrose (control) or 7 mmol/L sodium (control) beverage; b, indicates difference to 5% sucrose beverage or 15 mmol/L sodium beverage; c, indicates difference to 10% sucrose beverage. Statistical significance was accepted at $P < 0.05$. The vertical error bar in the top left corner represents the mean SD during the 4-h collection.

Figure 1

A



B




Figure 2




Figure 3




Figure 4





Figure 5

A**B**