



LJMU Research Online

Lake, MJ and Smith, G

The influence of sprint spike bending stiffness on sprinting

<http://researchonline.ljmu.ac.uk/id/eprint/3323/>

Article

Citation (please note it is advisable to refer to the publisher's version if you intend to cite from this work)


Lake, MJ and Smith, G (2016) The influence of sprint spike bending stiffness on sprinting. Footwear Science. ISSN 1942-4299

LJMU has developed **LJMU Research Online** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact researchonline@ljmu.ac.uk

<http://researchonline.ljmu.ac.uk/>



The influence of sprint spike bending stiffness on sprinting performance and metatarsophalangeal joint function

Journal:	<i>Footwear Science</i>
Manuscript ID	TFWS-2015-0139.R2
Manuscript Type:	Stiffness Thematic Issue
Keywords:	biomechanics, sprinting, bending, spikes, stiffness, longitudinal

SCHOLARONE™
Manuscripts

1. Background

The metatarsophalangeal joint (MTPJ) has been highlighted by several authors to have an important role in the energetics of the lower limb during sprint running (Bezodis, Salo & Trewartha, 2012; Smith, Lake, Lees, & Worsfold, 2012; Stefanyshyn & Nigg, 1997). In fact, this joint appears to be a net dissipater of energy during stance. It has been speculated by several authors that altering the mechanical properties of running footwear, in particular the longitudinal bending stiffness, is one mechanism by which the kinetics of the MTPJ may be altered (Smith, Lake, & Lees, 2014; Stefanyshyn & Fusco, 2004; Toon, Hopkinson & Kane, 2008) potentially leading towards slightly improved sprint performance. In vertical jumping, stiffer running shoes have been shown to significantly reduce MTPJ energy loss and consequently improve jump height (Stefanyshyn & Nigg, 2000). Similarly in running at 3.5m/s, an increase in the midsole longitudinal bending stiffness both significantly reduced the energy loss and at the same time increased positive work performed (Willwacher, Konig, Potthast, & Bruggemann, 2013).

For maximal sprint running, the effect of sprinting footwear bending stiffness on sprinting performance remains to be fully agreed between researchers, as there is some conflicting evidence in the literature regarding the benefit of wearing stiffer sprint shoes. Stefanyshyn and Fusco (2004) reported a decrease of 0.02s in 20-40 m sprint time for a group of 34 elite national sprinters when stiff carbon insoles were inserted into their running spikes (42 N·mm⁻¹ bending stiffness, measured using a three-point bending test). Conversely, Ding, Sterzing, Qin, and Cheung (2011) found no systematic influence of sprint spike stiffness upon 25 m acceleration performance for a group of young competitive athletes. When comparing barefoot sprinting to sprinting in sprint spikes, Smith et al. (2014) reported substantial changes in MTPJ function and potential improvements in performance-related parameters

1
2
3 due to wearing spikes. Sprint spikes appeared to have a clear localised effect on the function
4 of the MTPJ; increasing the work performed by lengthening the moment arm and likely
5 contributing to the energy produced during push-off. Stefanyshyn and Fusco (2004)
6 speculated that increasing the shoe bending stiffness would result in an anterior shift in the
7 point of application of the ground reaction force, thereby increasing the moment arm. This
8 would result in an increased MTPJ joint moment and energy production during late stance,
9 therefore enabling a more effective push-off. Willwacher et al. (2013) also observed that
10 increasing the midsole longitudinal bending stiffness seems to alter mechanical power
11 generation capacities of the MTPJ plantar flexing muscle tendon units by changing ground
12 reaction force leverage and decreasing plantar flexion angular velocity. Increasing the
13 longitudinal bending stiffness altered the lever arms at the MTPJ and ankle joint, changing
14 the gear ratio; the ratio of external GRF lever arm and internal lever arms of muscle tendon
15 units acting at a joint, and therefore potentially permitting the muscle tendon units to generate
16 higher joint moments (Willwacher et al., 2014). However, those authors pointed out that any
17 energy return at the metatarsophalangeal joint would require both the elastic midsole and the
18 toes to simultaneously perform a pronounced plantar flexion movement at the end of stance,
19 therefore energy would need to be transferred to the sprinter at the right place and the right
20 time.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 In elite sprinters, reported variation in lower limb joint kinetics within groups of sprinters
46 (Johnson & Buckley, 2001; Bezodis, Kerwin & Salo, 2008) suggests between-sprinter
47 differences do exist. This variation in technique is important to consider in the context of elite
48 sprinting, as even slight improvements in performance could affect finishing position within a
49 sprint race (Bezodis, Salo, & Trewartha, 2014). Similarly, individual athletes may elicit
50 different biomechanical and performance responses to changes in footwear, suggesting that
51
52
53
54
55
56
57
58
59
60

1
2
3 shoe selection is specific to the functional requirements of individuals (Toon, Williams,
4
5 Hopkinson & Kane, 2009) and might (among others) depend upon their ankle plantarflexor
6
7 strength, force-length and force-velocity characteristics (Stefanyshyn & Fusco, 2004).
8
9 Differences in responses to running shoes of varying stiffness coincide with runner's
10
11 preferences and subjective perception (Kleindienst, Michel, & Krabbe 2005). It has been
12
13 suggested that the only way to detect differences that exist between shoes is to employ a
14
15 single-subject analysis (Bates, 1996). Furthermore, individual analysis has been advocated
16
17 for future research to detecting differences between types of running shoes, based upon high
18
19 variability of ground reaction forces reported in male and female groups of runners (Logan et
20
21 al., 2010). Single subject analyses have been advocated in the research when a task could
22
23 potentially elicit different responses / strategies in individuals and when the group mean does
24
25 not necessarily describe well any individual subject response (Revill, Perry, Edwards, &
26
27 Dickey, 2008).
28
29
30
31
32
33
34

35 Therefore the aim of this study was to first investigate the effect of sprint spike bending
36
37 stiffness on sprinting performance and MTPJ motion in a group of sprinters. This
38
39 experimental design was supplemented by a second investigation using the same footwear
40
41 modifications during sprinting but utilising a single subject design and improved
42
43 biomechanical data collection and analysis (Smith et al., 2012). It was hypothesised that
44
45 group findings may mask some significant individual changes in foot function and sprint
46
47 performance with sprint shoe stiffness.
48
49

50 51 **2. Methods**

52
53 This research comprised of two studies, a group study, followed by an individual case-study.
54
55

56 57 *Experimental footwear*

1
2
3 Commercially available sprint spikes (Puma Complete Theseus II) in two sizes (38.5 and 43
4 EUR) were provided by Puma SE, Germany and custom made insoles of varying longitudinal
5 bending stiffnesses, made combining different layers of carbon and glass fibre, were inserted
6 into the identical shoes. Additionally, Asics Hyper Sprint shoes of same sizes were used,
7 serving as additional controls (A-control). The bending stiffness of the five sprint spike
8 conditions of each size were measured mechanically: A-control and four Puma test
9 conditions; P-control (no insole), P-low, P-medium, P-high (added stiffnesses with insoles).
10 A two point bending test was used (Krumm, Schwanitz & Oldenwald, 2012; Smith et al.,
11 2014) with 40 mm of bending at a constant velocity of 10 mm/s. A Servo hydraulic material
12 testing machine was used (Zwick GmbH & Co. KG, Ulm, Germany, stroke 100 mm, load
13 max. 10 kN) with a LVDT position sensor and a 10 kN load cell (Huppert GmbH Prüf- und
14 Messtechnik, Herrenberg, Germany) with the sprint spikes placed in a mould and secured
15 with a metal clamp on the forefoot. The average stiffness for a deformation of 0-40 mm
16 dorsiflexion at the natural bending line was measured for three trials and then averaged per
17 left and right shoe (Table 1).
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 *Group Study*

37
38
39 Twelve competitive athletes (club / regional level) participated in the group study; six male
40 (mean age 22.0 ± 3.6 years, mean height 182.7 ± 4.5 cm, mean mass $73.8 \text{ kg} \pm 4.4 \text{ kg}$, mean
41 100 m personal best 11.3 ± 0.3 s). and six female (mean age 21.8 ± 4.8 years, mean height
42 167.0 ± 6.2 cm, mean mass $60.2 \pm 4.2 \text{ kg}$, mean 100 m personal best 12.4 ± 0.3 s). Informed
43 written consent was obtained in accordance with the Ethics Committee of Technische
44 Universität Chemnitz. All participants fitted shoe size 43 or 38.5.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 After their own warm up and two practice runs at 80% and 100% of their maximum velocity,
4 sprinters performed eight maximal 40 m sprints from a standing start, along an indoor 100 m
5 athletics track at Technische Universität Chemnitz. This consisted of two sprint trials in each
6 Puma footwear condition (P-control, P-low, P-medium, and P-high) and athletes were asked
7 to sprint maximally. The order of sprint spikes was randomised and counter-balanced and
8 participants were blind to the footwear condition. Eight minutes rest between each maximal
9 sprint was given, based upon pilot work examining fatigue over multiple sprint trials (Figure
10 1). 10 m split time was taken from 30 - 40 m using single beam timing cells. A high speed
11 video camera (Exilim Pro Ex-F1, 600 Hz sample rate, 1/1000 s shutter speed) captured one
12 foot contact 15 m into the sprint (acceleration phase). High speed videos were analysed in
13 Quintic Biomechanics (Quintic Consultancy Ltd.) where the following points were manually
14 digitised: the heel; the distal end of the hallux and the medial side of the first metatarsal head
15 (MTH1). For two-dimensional analysis, a marker on MTH1 or MTH2 provides a better
16 representation of MTPJ motion than a marker on MTH5 (see Smith et al., 2012). The MTPJ
17 angle was therefore defined as the angle between the forefoot (MTH1 – hallux) and rearfoot
18 (MTH1 - heel) segments. The raw kinematic data was smoothed using a 4th order Butterworth
19 low pass digital filter with cut off frequencies of 60 Hz for the MTPJ angle and 30 Hz for the
20 angular velocity.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 *Statistical analysis*

44
45
46 A one-way repeated measures ANOVA was conducted to evaluate the intervention of sprint
47 spikes stiffness (mean of two trials per condition) on sprint time, MTPJ range of motion,
48 MTPJ maximum flexion and MTPJ peak angular velocities, both plantarflexion and
49 dorsiflexion.
50
51
52
53
54

55 *Individual case study*

1
2
3 This aspect of the research work utilised a novel experimental design that aimed to tease out
4 the likely small differences in sprint performance and lower limb function when sprinting in
5 shoes of different stiffness. This part of the research work represents both an evolution to the
6 study design and improvements to the biomechanical measurements taken, to gain further
7 insight into individual responses in sprinting performance and MTPJ function to modified
8 bending stiffness of sprint spikes. This study took place at a different testing location to
9 enable the Biomechanical analysis.
10
11

12
13
14
15
16
17
18
19 Two additional trained, competitive sprinters took part in the individual study; one female
20 100 m sprinter / hurdler (aged 28 years, height 179cm, mass 68 kg, shoe size 38.5, 100 m
21 sprint time 12.2 s) and one male 400 m specialist (aged 28 years, height 182cm, mass 73 kg,
22 shoe size 43, 400 m best time 50.8 s). These two participants were chosen for the individual
23 study as they were also experienced, well-trained track athletes, whose mean sprinting
24 velocities at 35 m reflected the mean 30 - 40 m sprinting velocities of the group study
25 (within one standard deviation). Informed written consent was obtained in accordance with
26 the Ethics Committee of Liverpool John Moores University.
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 Each sprinter attended the lab three times, each time to test a different sprint spike stiffness
42 (P-low, P-medium or P-high) condition in relation to A-control, therefore three stiffness
43 conditions were tested in relation to control (Figure 1). The design included a balanced,
44 presentation of the control shoe condition (A control) and a test shoe condition (P-low, P-
45 medium or P-high) which minimises the error sources associated with presenting different
46 test stimuli which are separated by a period of time (Gescheider, 1997). In each testing
47 session, six sprinting trials per shoe were recorded, with the sprinter swapping shoes between
48 each trial, e.g. A-control followed by P-low, repeated six times. Therefore, a total of twelve
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 maximal sprints were collected per testing sessions, this number was based upon pilot testing
4
5 of the number of sprints practically possible. The sprints were performed on a 70 m indoor
6
7 sprint track at Liverpool John Moores University and the athletes accelerated for
8
9 approximately 35 m before data collection. Athletes were instructed to sprint maximally to 40
10
11 m. A single left foot ground contact in the middle of a force platform (Kistler model 9287B,
12
13 sampling at 1000 Hz) at approximately 35 m was used for analysis. A customized starting
14
15 mark was used to aid the athlete in striking the force plate without the need to alter their
16
17 stride pattern prior to force plate contact. Single beam timing cells were located 2.5 m either
18
19 side of the force platform, therefore recording sprint times over 5 m as the athletes crossed
20
21 the force platform. The athletes were still accelerating at this point.
22
23
24
25
26
27

28
29 Both participants underwent two DEXA scans to optimise the location of markers on the
30
31 metatarsal heads (Smith et al., 2014). Kinematic data were collected using a 6 camera system
32
33 (Pro-Reflex MCU 1000, Qualisys Inc., Sweden) sampling at 1000 Hz. Data were processed
34
35 using Visual3D (C-Motion, Inc.) and a three-segment foot model was used for the kinematic
36
37 analysis and kinetic analysis as detailed in Smith et al. (2012). In brief, the five MTPJ's were
38
39 considered as a single joint rotating about an axis oblique to the sagittal plane defined by
40
41 markers (11 mm diameter) on MTH1 and 5. Holes were cut out in the spikes for markers to
42
43 be located on MTH1, MTH5 and MTH2. MTH2 was a virtual marker identified using a 3D
44
45 pointer device (C-motion Inc., Canada). When the sprint spikes were swapped, a new
46
47 standing calibration and pointer trial was performed. Joint positional and force data were
48
49 smoothed using a fourth-order low pass Butterworth filter with a cut-off frequency (F_c) of
50
51 100 Hz. This was necessary to preserve the high frequency aspects of the signal previously
52
53 identified and important for determining the presence of any positive joint power during the
54
55 push-off phase (Smith et al. 2012). Force platform point of force application (PFA)
56
57
58
59
60

1
2
3 measurements were determined above a vertical force threshold of 100 N for initial landing
4
5 and 50 N at the end of ground contact, based upon visual inspection of PFA curves, as errors
6
7 were greater at the start of foot contact where higher loading rates were experienced. Below
8
9 these thresholds the PFA was distorted and in a position outside of the forefoot, due to low
10
11 loading on the force platform. MTPJ joint moments and powers were calculated when the
12
13 centre of pressure moved distally to its oblique axis during sprinting ground contact.
14
15

16 17 *Statistical analysis*

18
19 For each sprinter's within-subject statistical analysis, the Model Statistic was used. This
20
21 approach was chosen to replicate previous single-subject studies assessing performance of
22
23 sports shoes (Dufek & Bates, 1991). The Model Statistic is the single-subject equivalent of an
24
25 independent t-test (Bates, James, & Dufek 2004) as using a repeated-measures approach in a
26
27 single subject design is inappropriate, therefore the more appropriate approach is to treat trial
28
29 values as independent measures and use the corresponding independent test procedure (Bates,
30
31 1996). The mean absolute differences between two conditions (for example MTPJ Moment in
32
33 a stiff shoe minus MTPJ moment in the control condition) is calculated and compared to a
34
35 critical difference. The critical difference is the product of the standard deviation (of all trials
36
37 in each footwear condition) and a critical value based upon sample (trial) size from Bates et
38
39 al., (2004). The critical value used in this study was 1.2408 for six trial comparisons (twelve
40
41 trials in total) and $\alpha = 0.05$ level of significance. The interpretation of a mean absolute
42
43 difference greater than the critical difference is that the difference is due to sampling error at
44
45 a probability of α , therefore the null hypothesis is rejected. The single-subject, repeated trials
46
47 design also accounts for any learning or fatigue during the testing session, as every trial in a
48
49 stiffness condition is compared to the control (Figure 1).
50
51
52
53
54

55 56 **3. Results**

57 58 *Group study*

1
2
3 Thus, bending stiffness values of the sprint spikes used in this study indicate that, by adding
4 the insoles, bending stiffness was increased considerably in comparison to the commercial
5 sprint spike conditions used, P-high had almost three times greater stiffness (273%) than the
6 A-control shoe (Table 1). Furthermore, four additional commercially available sprint spikes
7 were mechanically tested and had average stiffnesses ranging from 190 ± 5.3 N/m to $256 \pm$
8 23.7 N/m.
9
10
11
12
13
14
15
16
17
18
19

20 Average 30 m to 40 m sprint time for all trials was 1.18 s (± 0.08 s), corresponding to a mean
21 velocity of 8.50 m/s (± 0.57 m/s). There were no significant differences ($p < 0.05$) in mean
22 sprinting velocity across the shoe stiffness conditions (Figure 2) ($p = 0.634$, $F = 0.427$).
23
24
25
26
27
28
29

30 Individual differences existed between sprinters with some sprinters appearing to perform
31 slightly better in P-control and others possibly performing slightly better with stiff insoles
32 (Figure 3). Observations of individual performance differences between shoe stiffness
33 conditions did not appear to be related to overall sprinting ability. Both faster and slower
34 sprinters demonstrated either increasing or decreasing sprint performances with increasing
35 shoe bending stiffness (Figure 3).
36
37
38
39
40
41
42
43
44
45
46

47 There were no significant differences in the metatarsophalangeal range of motion during
48 stance with mean values of 40.5° ($\pm 3.0^\circ$), 38.1° ($\pm 4.5^\circ$), 38.3° ($\pm 2.8^\circ$) and 37.5° (± 4.7
49 $^\circ$) for P-control, P-low, P-medium and P-high respectively ($p = 0.277$, $F = 1.510$). Nor were
50 there any statistically significant differences in maximum dorsiflexion ($p = 0.600$, $F = 0.654$)
51 or plantarflexion ($p = 0.819$, $F = 0.109$) angular velocities across the four conditions.
52
53
54
55
56
57
58
59
60

Individual case study

The female participant achieved highest mean sprinting velocities in the P-medium and P-high, however the only significant difference, to A-control, was found for P-high, which resulted in a 2% increase in measured velocity (Table 3). The male participant achieved highest mean sprinting velocities in P-medium shoe, however there were no significant differences in velocities between any of the P stiffness conditions and A-control (Table 2). Tables 2 and 3 show individualised differences in MTPJ kinematics and kinetics between the conditions.

The female participant exhibited significantly higher resultant MTPJ moments in all three P stiffness conditions, in comparison to A-control (Table 3). In P-high the joint moment was increased by approximately 8.9 N·m (a 15 % increase). P-low and P-high significantly increased power production during the push-off phase of sprinting, although there were no favourable reductions in energy loss with the stiffened shoe conditions. Similarly, the male participant demonstrated significant differences in work performed at the MTPJ with significantly higher joint moments in all P stiffness conditions, in comparison to A-control, with P-high condition resulting in the highest joint moment (Figure 4). The MTPJ was a net absorber of energy during stance with up to 42.0 J lost during the energy absorption phase in the A-control (Table 2). All three P shoe conditions significantly reduced the amount of energy lost, compared to A-control. The largest reduction was in P-medium with 8.2 J (23%) difference in energy loss compared to A-control. A significantly increased positive energy was created during the push-off phase at the end of stance in P-low, producing 3.3 J energy, compared to 2.2 J energy in the A-control.

4. Discussion and Conclusion

1
2
3 This study found that increased sprint shoe bending stiffness was generally not linked with
4 improved sprinting performance for a group of trained sprinters. The small improvement in
5 performance reported by Stefanyshyn and Fusco (2004) may be due to a larger number of
6 subjects used (34 sprinters) and a more homogeneous group of older elite athletes. However,
7 the authors argued that performance enhancements of 0.7% are quite large, and therefore
8 worthwhile, in elite sprinting, possibly making a difference in finishing position in a race.
9 The lack of group mean shoe differences seen in the group study may be the result of high
10 variability in individual results. The number of subjects ($n = 12$) and trials used in this study
11 was not dissimilar to, or greater than previous biomechanical studies using trained sprinters
12 (Bezodis et al., 2008; Ding et al., 2011; Gittoes & Wilson, 2010; Johnson & Buckley, 2001;
13 Stefanyshyn & Nigg, 1997), although it is acknowledged that the sprinters in this study were
14 younger and not world-class. The results from this study do, however, agree with the
15 suggestion that the stiffness each athlete requires to produce maximal performance is subject-
16 specific (Stefanyshyn & Fusco, 2004). Therefore, it is suggested that an athlete's optimal
17 sprint shoe stiffness is individualised and likely dependent upon their perceptions and
18 previous experiences (Kleindenst, Michel & Krabbe, 2005) forefoot structure or stiffness
19 (Oleson, Adler & Goldsmith, 2005), and arch stiffness (Butler, Davis & Hamill, 2006).

20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44 In the individual case studies, the female participant demonstrated significantly improved
45 sprinting performance in P-high stiffness condition, whilst improvements in sprinting speed
46 for the male participant were not significant. For both participants, stiffer sprint spikes did
47 appear to have a controlling effect over the MTPJ kinematics, significantly reducing the
48 range of motion and also reducing the extent and rate of MTPJ dorsiflexion during stance and
49 plantarflexion after push-off. This reaffirms, on an individual level, the restrictive influence
50 that stiffened footwear seems to have on the normal bending of the MTPJ during stance,
51
52
53
54
55
56
57
58
59
60

possibly influencing the effectiveness of the Windlass mechanism (Hicks, 1954), although this would likely be dependent upon the individual's foot dynamic stiffness. However, the controlling effect over the behaviour of the foot, does not appear to adversely affect energy production of the joint (Smith et al., 2014) and both athletes were able to produce energy within the foot/shoe complex during the push-off phase of sprinting. This is in agreement with Lin et al., (2013) who demonstrated that for walking, carbon fibre insoles effectively reduced the windlass effect, but also facilitated storage of energy that can be used for efficient gait propulsion without needing to dorsiflex the MTPJ to a great extent.

The results of this study suggest that stiffer sprint spikes may increase the loading applied on the central forefoot and may also facilitate greater anterior progression of the centre of pressure, as evidenced by a greater moment arm, to enable increased propulsion during push off. Values for MTPJ moments and a substantial net loss of energy at this joint are similar to those reported in other studies for sprinting (Smith et al., 2014; Stefanyshyn & Nigg, 1997; Toon et al., 2009). It is clear that the longitudinal stiffness of sprint spikes have a clear localised effect on the MPTJ, increasing the work performed on the joint, lengthening the moment arm and producing small amounts of energy during push-off, in agreement with Smith et al., 2014. However, it is also likely that the range of motion, joint moment and resultant power at the ankle joint will also be altered by forefoot bending stiffness (Chen et al. 2014). Wilwacher et al., (2014) demonstrated increasing the longitudinal bending stiffness in running shoes led to a significant anterior shift of all lower limb lever arms, this effect greater at the more distal joints In particular, higher ankle joint moments might be beneficial for increasing storage and return of energy in the Achilles Tendon, particularly in elite sprinters who are likely to have higher strength potential of the plantar flexor muscle tendon units.

1
2
3
4
5
6 It is impossible to compare the bending stiffness to other studies, due to different methods of
7
8 quantifying longitudinal bending stiffness, plus the unknown stiffness of sprinters' own
9
10 sprinting shoes. However, in this current study, the mechanical stiffness of the whole sprint
11
12 spike system was determined mechanically and every stiffness condition was quantified and
13
14 compared to a range of commercially available sprint spikes. It is also acknowledged that the
15
16 testing protocol was modified from the group study to the individual case-study. However,
17
18 this was done so in order to improve both the study design (number of testing session trials
19
20 and conditions), and biomechanical measurements (MTPJ kinetics) possible for exploring
21
22 individual responses to sprint spike stiffness in greater detail.
23
24

25
26 The high bending stiffness of the sprint spikes may allow the athletes to push off but still
27
28 achieve substantial rigidity from the foot and shoe as a system. This concurs with the findings
29
30 of Willwacher et al. (2013) for lower-velocity running; an increase in shoe stiffness around
31
32 the MTPJ provides better working conditions for plantarflexing muscle tendon units,
33
34 increasing the push-off duration. Participant one exhibited higher ranges of motion than
35
36 participant two, indicating differences between the participants in their natural range of
37
38 motion and flexibility within the foot and differing inherent foot stiffness or foot structures
39
40 (e.g. high / low arch, plantar fascia stiffness). This is concurrent with the wide range of arch
41
42 stiffness reported in high arch (1993.8 ± 1112.1 kg/arch height index) and low arch ($788.0 \pm$
43
44 407.0 kg/arch height index) runners (Butler et al., 2006) Participant two experienced larger
45
46 reductions in angular motion and peak angular velocities with the stiffer sprint spikes,
47
48 suggesting greater responses to shoe stiffness. Both participants demonstrated increased joint
49
50 moments with increased stiffness of sprint spike. Energy production during the last 10
51
52 milliseconds of stance was also increased by a stiffer sprint shoe.
53
54
55
56
57
58
59
60

1
2
3
4
5
6 The two participants in this study exhibited different responses to the footwear conditions,
7
8 this may be due to anatomical differences and neurological responses (performance
9
10 strategies). The participant's response has been suggested to depend on their recognition /
11
12 perception of the perturbation, which in turn will be a function of the individual's past
13
14 experiences (Bates, 1996). Response patterns vary along a continuum (from a Newtonian
15
16 response whereby the perturbation is completely ignored to a fully accommodating response).
17
18 Individual performance strategies are likely to be the norm rather than the exception, due to
19
20 the complexity of the human machine and its numerous associated functional degrees, along
21
22 with the different experiences, perceptions and expectations of the participants (Bates, 1996).
23
24
25
26
27
28

29 The individual responses exhibited in this study suggest that sprinting performance may be
30
31 improved by implementation of relevant shoe mechanical characteristics. It appears that shoe
32
33 selection may be specific to the functional requirements of individual athletes and athletes
34
35 will likely respond differently to footwear modifications. Future work should identify key
36
37 anatomical or biomechanical factors that influence foot function during maximal sprinting.
38
39 For example, the influence of the natural stiffness of the human forefoot, and the strength and
40
41 velocity characteristics of the ankle plantarflexors and toe flexors, are morphological factors
42
43 which may influence foot function during sprinting and may dictate appropriate shoe
44
45 selection. Besides changes to the overall longitudinal bending stiffness, other mechanical
46
47 characteristics of sprint spike design, such as the toe spring angle warrant future
48
49 investigation. The importance of appropriately modelling the position and orientation of the
50
51 MTPJ has been previously established (Smith et al., 2012), therefore the location of the
52
53
54
55
56
57
58
59
60

1
2
3 natural bending line of the sprint spikes in relation to the anatomical joint axis will likely
4
5 affect the resultant motion at the foot.
6
7

8 Overall, the findings suggest that high variability in individual responses to sprint shoe
9
10 bending stiffness manipulations can be masked by group mean findings and a single-subject
11
12 approach appears warranted to further investigate whether shoe bending stiffness can be
13
14 optimised to improve individual sprint performance.
15
16

17 **Acknowledgments**

18
19
20 Authors are grateful to the International Society of Biomechanics for the award of a Student
21
22 Travel Grant to support this study.
23
24

25 **References**

26
27
28 Bates, B.T. (1996). Single-subject methodology: an alternative approach. *Medicine and*
29
30 *Science in Sports and Exercise*, 28, 631-638.
31
32

33 Bates, B.T., James, C.R., & Dufek, J. (2004). Single subject Analysis. In N.Stergiou (Ed.),
34
35 *Innovative Analysis of Human Movement* (pp. 3 - 28). Human Kinetics: Champaign IL.
36
37

38 Bezodis, I.N., Kerwin, D.G., & Salo, A.I.T. (2008). Lower-Limb Mechanics during the
39
40 Support Phase of Maximal-Velocity Sprint Running. *Medicine and Science in Sports and*
41
42 *Exercise*, 40, 707-715.
43
44

45 Bezodis, N.E., Salo, A. I.T., & Trewartha, G. (2012). Modelling the Stance Leg in 2D
46
47 Analyses of Sprinting: Inclusion of the MTP Joint Affects Joint Kinetics. *Journal of Applied*
48
49 *Biomechanics*, 28, 222-227.
50
51

52 Bezodis, N.E., Salo, A. I.T., & Trewartha, G. (2014). Lower limb joint kinetics during the
53
54 stance phase in athletics sprinting: three elite athlete case studies. *Journal of Sports Sciences*,
55
56 32, 738-746.
57
58
59
60

1
2
3 Bisseling, R.W., & Hof, L.A. (2006). Handling of impact forces in inverse dynamics. *Journal*
4 *of Biomechanics*, 39, 2438-2444.
5
6

7
8 Butler, R.J., Davis, I.S., & Hamill, J. (2006). Interaction of Arch Type and Footwear on
9 Running Mechanics. *American Journal of Sports Medicine*, 34, 1998-2005.
10
11

12 Chen, C-H., Tu, K-H., Liu, C., & Shiang T-Y. (2014). Effects of forefoot bending elasticity of
13 running shoes on gait and running performance. *Human Movement Science*, 38, 163-172.
14
15

16
17 Ding, R., Sterzing, T., L. Quin, T., & Cheung, J. (2011). Effect of metatarsal-phalangeal joint
18 and sprint spike stiffness on sprint spike performance. *Footwear Science*, 3:sup1, S41-S43.
19
20

21
22 Dufek, J.S., & Bates, B.T. (1991). Dynamic performance assessment of selected sports shoe
23 on impact forces. *Medicine and Science in Sports and Exercise*, 23, 1062-1067.
24
25

26
27 Gescheider, G. A. (1997). *Psychophysics: The fundamentals* (4th ed.). New Jersey: Erlbaum.
28
29

30
31 Gittoes, M.J.R., & Wilson, C. (2010) Intralimb joint Coordination Patterns of the Lower
32 Extremity in Maximal Velocity Phase Sprint Running. *Journal of Applied Biomechanics*, 2,
33 188 – 195.
34
35

36
37
38 Hicks, J.H. (1954). The mechanics of the foot II. The plantar aponeurosis and the arch.
39 *Journal of Anatomy*, 87, 345-357.
40
41

42
43 Johnson, M.D., & Buckley, J.G. (2001). Muscle power patterns in the mid-acceleration phase
44 of sprinting. *Journal of Sports Sciences*, 19, 263-272.
45
46

47
48 Kleindienst, F.I., Michel, K.J., & Krabbe, B. (2005). Influence of midsole bending stiffness
49 on the metatarsophalangeal joint based on kinematic and kinetic data during running.
50 *Proceedings of the 7th ISB footwear biomechanics symposium*, Cleveland, Ohio.
51
52
53
54
55
56
57
58
59
60

1
2
3 Krumm, D., Schwanitz, S., & Oldenwald, S. (2012). Development and reliability
4
5 quantification of a novel test set-up for measuring footwear stiffness. *Sports Engineering*, 16,
6
7 13-19.

8
9
10 Lin, S-H., Chen, C., Tang, S., Wong, A., Hsieh, J-H., & Chen, W-P. (2013). Changes in
11
12 windlass effect in response to different shoe and insole designs during walking. *Gait &*
13
14 *Posture*, 37, 235-241.

15
16
17 Logan, S., Hunter, I., Hopkins, J.T., Feland, J.B. & Parcell, A. (2010). Ground Reaction
18
19 Force Differences between Running Shoes, racing Flats, and Distance Spikes in Runners.
20
21 *Journal of Sports Science and Medicine*, 9, 147-153.

22
23
24 Oleson, M., Adler, D., & Goldsmith, P. (2005). A comparison of forefoot stiffness in running
25
26 and running bending stiffness. *Journal of Biomechanics*, 38, 1886-1894.

27
28
29 Revill, A.I., Perry, S.D., Edwards, A.M., & Dickey, J.P. (2008). Variability of the impact
30
31 transient during repeated barefoot walking trials. *Journal of Biomechanics*, 41, 926-930.

32
33
34 Smith, G.E, Lake, M.L., Lees, A., & Worsfold P. (2012). Measurement procedures effect the
35
36 interpretation of metatarsophalangeal joint function during accelerated sprinting. *Journal of*
37
38 *Sports Sciences*, 30, 1521-7.

39
40
41 Smith, G.E., Lake M.L., & Lees, A. (2014). Metatarsophalangeal joint function during
42
43 sprinting: A comparison of barefoot and sprint spike shod foot conditions. *Journal of Applied*
44
45 *Biomechanics*, 30, 206-212.

46
47
48 Stefanyshyn, D.J., & Fusco, C. (2004) Increased shoe bending stiffness increases sprint
49
50 performance. *Sports Biomechanics*, 3, 55-66.

51
52
53 Stefanyhysn, D.J., & Nigg, B.M. (1997). Mechanical energy contribution of the
54
55 metatarsophalangeal joint to running and sprinting. *Journal of Biomechanics*, 20, 1081-1085.
56
57
58
59
60

1
2
3 Stefanyshyn, D.J., & Nigg, B.N. (2000). Influence of midsole bending stiffness on joint
4 energy and jump height performance. *Medicine and Science in Sports and Exercise*, 32, 471-
5 476.
6
7

8
9
10 Toon, D., Hopkinson, N., & Caine, M. (2008). The effect of shoe bending stiffness on
11 predictors of sprint performance. In A. Subic, F. K., Fuss, & S. Ujihashi (Eds.), *The Impact of*
12 *Technology on Sport II* (pp. 643-648). Taylor & Francis.
13
14

15
16
17 Toon, D., Williams, B., Hopkinson, N., & Caine, M. (2009). A comparison of barefoot and
18 sprint spike conditions in sprinting. *Journal of Sports Engineering and Technology;*
19 *Proceedings of the Institute of Mechanical Engineers, Part P*, 223, 77-87.
20
21

22
23
24 Willwacher, S., Konig, M., Potthast, W., & Bruggemann, G-P. (2013). Does specific
25 footwear facilitate energy storage and return at the metatarsophalangeal joint in running?
26 *Journal of Applied Biomechanics*, 29, 583-592.
27
28

29
30
31
32 Willwacher, S., Konig, M., Braunstein, B., Goldmann, J-P., & Bruggemann, G-P. (2014).
33 The gearing function of running shoe longitudinal bending stiffness. *Gait & Posture*, 40, 386-
34 390.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Schematic demonstrating study design and protocol for the Group Study and Individual Case Study




Figure 2. Mean (SD) sprinting velocity for twelve sprinting subjects performing two sprints in four sprint spike stiffness conditions.




Figure 3. 10 m split (30 – 40m) sprint time for twelve sprinting participants performing two sprints each in four sprint spike bending stiffness conditions. Each line represents a different sprinter.




Figure 4. Mean resultant MTPJ moment during the stance phase of sprinting for male sprinter in the individual case study, for condition A-control; solid line, and P-medium; dashed line. Peak moment was significantly ($p < 0.05$) different between the two bending stiffness conditions.

Table 1. Average stiffness (and stiffness normalised to the Puma no insole condition) of the sprint spike conditions used in the study, left and right shoes.

Shoe condition	Shoe size 38.5 Average Mechanical stiffness (N/m) of left and right shoe and Normalised stiffness to condition 0 (%)	Shoe size 43 Average Mechanical stiffness (N/m) of left and right shoe and Normalised stiffness to condition 0(%)
Puma no insole	254.6 ± 22.7 (100%)	297.4 ± 7.6 (100%)
Puma low stiffness insole (low)	314.6 ± 7.8 (124%)	343.1 ± 6.0 (115%)
Puma medium stiffness insole (medium)	367.0 ± 19.7 (144%)	408.6 ± 10.5 (± 137%)
Puma high stiffness insole (high)	501.7 ± 43.6 (197%)	472.1 ± 31.6 (159%)
Asics Baseline	183.66 ± 6.25 (72%)	197.9 ± 29.6 (67%)

Table 2. Male sprinter's mean sprinting velocity and selected MTPJ kinematic and kinetic variables for low, medium and high stiffness conditions to corresponding baseline conditions (Asics baseline). * denotes a significant difference to the corresponding baseline measure ($p < 0.05$).

Stiffness	Asics baseline	Low	Asics baseline	Medium	Asics baseline	High
Speed (m/s)	7.66 (± 0.22)	7.60 (± 0.15)	8.05 (± 0.27)	8.12 (± 0.27)	7.94 (± 0.21)	7.90 (± 0.24)
MTPJ Range of Motion ($^{\circ}$)	37.1 (± 1.4)	31.6* (± 1.5)	37.8 (± 1.2)	28.5* (± 2.0)	36.5 (± 1.6)	30.3* (2.5)
Peak Dorsiflexion angular velocity ($^{\circ}/s$)	844.2 (± 49.1)	703.3* (± 84.2)	773.6 (± 74.3)	534.4* (± 46.4)	783.8 (± 44.8)	697.2* (± 68.5)
Peak Plantarflexion angular velocity ($^{\circ}/s$)	-2359.0 (± 95.0)	-2146.8* (± 107.0)	-2507.4 (± 175.4)	-2150.2* (± 187.9)	-2288.0 (± 392.7)	-2120.2* (± 331.5)
Peak vertical force Fz (N)	2261.9 (± 60.0)	2194.5* (± 79.5)	2271.7 (± 36.9)	2345.6* (± 72.1)	2253.2 (± 98.6)	2324.9 (± 89.1)
Peak plantar flexor moment (N·m)	83.1 (± 2.4)	87.6* (± 2.9)	72.1 (± 5.0)	79.7* (± 3.3)	75.8 (± 4.6)	87.6* (± 6.6)
Peak Positive Power (W) generated during plantarflexion	220.3 (± 23.9)	292.3 (± 73.5)	241.9 (± 24.2)	241.8 (± 22.0)	265.5 (± 52.3)	386.5* (± 128.5)
Peak Negative Power (W) generated during dorsiflexion	-1053.7 (± 78.1)	-976.2* (± 97.6)	-963.8 (± 102.4)	-678.2* (± 88.2)	-927.2 (± 78.2)	-916.2 (± 121.0)
Total Energy generated (J) during plantarflexion	0.6 (± 0.4)	0.4 (± 0.2)	0.2 (± 0.2)	0.2 (± 0.2)	0.5 (± 0.2)	0.8 (± 0.6)
Total Energy absorbed (J) during dorsiflexion	-42.0 (± 2.3)	-35.8* (± 3.0)	-35.0 (± 4.1)	-26.3* (± 1.3)	-37.8 (± 2.0)	-34.2* (± 3.3)
Total energy generated (J) during push-off	2.2 (± 0.3)	3.3* (± 0.6)	2.4 (± 0.4)	2.7 (± 0.4)	2.8 (± 0.7)	3.2 (± 0.6)

Table 3. Female sprinter's mean sprinting velocity and selected MTPJ kinematic and kinetic variables for low, medium and high stiffness conditions and corresponding baseline conditions (Asics baseline). * denotes a significant difference to the corresponding baseline measure ($p < 0.05$).

Stiffness	Asics baseline	Low	Asics baseline	Medium	Asics baseline	High
Speed (m/s)	8.15 (± 0.23)	8.14 (± 0.14)	8.07 (± 0.18)	8.20 (± 0.08)	8.04 (± 0.14)	8.20* (± 0.09)
Metatarsophalangeal Range of Motion ($^{\circ}$)	37.9 (± 3.8)	42.3* (± 3.1)	41.1 (± 2.5)	39.2 (± 1.2)	46.8 (± 2.0)	41.3* (3.0)
Peak Dorsiflexion angular velocity ($^{\circ}/s$)	1182.7 (± 314.4)	1228.3 (± 156.7)	1128.2 (± 158.1)	1068.5 (± 184.8)	1334.2 (± 261.3)	1197.8 (± 66.0)
Peak Plantarflexion angular velocity ($^{\circ}/s$)	-3071.0 (± 143.1)	-2729.3* (± 209.7)	-3472.0 (± 272.9)	-3008.5* (± 280.1)	-3080.8 (± 48.7)	-2476.5* (± 170.2)
Peak vertical force Fz (N)	2057.8 (± 36.3)	2144.0* (± 36.7)	2083.2 (± 39.1)	2136.1* (± 36.6)	2037.2 (± 34.8)	2078.1* (± 29.3)
Peak Plantarflexor moment (N.m)	53.7 (± 5.6)	64.0* (± 5.9)	56.0 (± 4.4)	72.7* (± 7.8)	58.7 (± 9.1)	67.6* (± 11.5)
Peak Positive Power (W) generated during plantarflexion	201.2 (± 100.0)	292.3 (± 73.5)	183.7 (± 81.0)	253.3 (± 33.9)	333.9 (± 148.9)	266.5 (± 57.7)
Peak Negative Power (W) generated during dorsiflexion	-691.7 (± 61.2)	-867.0 (± 55.0)	-777.4 (± 90.9)	-855.1 (± 90.6)	-1000.2 (± 169.2)	-927.9 (± 50.0)
Total Energy generated (J) during plantarflexion	1.4 (± 1.1)	0.5* (± 0.2)	1.3 (± 1.0)	0.9 (± 0.5)	2.0 (± 1.5)	1.3 (± 0.9)
Total Energy absorbed (J) during dorsiflexion	-30.6 (± 2.3)	-33.3 (± 2.0)	-31.2 (± 2.8)	-36.4* (± 3.9)	-34.1 (± 4.3)	-36.9 (± 6.6)
Total energy generated (J) during push-off	0.3 (± 0.1)	2.3* (± 0.6)	1.6 (± 0.7)	2.1 (± 0.4)	0.9 (± 0.4)	2.1* (± 0.5)