



## LJMU Research Online

**Kooreman, Z, Giraldeau, G, Finocchiaro, G, Kobayashi, Y, Wheeler, M, Perez, M, Moneghetti, K, Oxborough, D, George, KP, Myers, J, Ashley, E and Haddad, F**

**Athletic Remodeling in Female College Athletes, the "Morganroth Hypothesis" Revisited.**

<http://researchonline.ljmu.ac.uk/id/eprint/8372/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Kooreman, Z, Giraldeau, G, Finocchiaro, G, Kobayashi, Y, Wheeler, M, Perez, M, Moneghetti, K, Oxborough, D, George, KP, Myers, J, Ashley, E and Haddad, F (2018) Athletic Remodeling in Female College Athletes, the "Morganroth Hypothesis" Revisited. Clinical Journal of Sport Medicine.**

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk/) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

## **Athletic Remodeling in College Female Athletes, the “Morganroth Hypothesis” Revisited**

Zoe Kooreman, MD<sup>1,2</sup>, Geneviève Giraldeau, MD<sup>1,2</sup>, Gherardo Finocchiaro, MD<sup>1,2</sup>, Yukari Kobayashi, MD<sup>1,2</sup>, Matthew Wheeler, MD<sup>1,2</sup>, Marco Perez, MD<sup>1,2</sup>, Kegan Moneghetti, MD, David Oxborough PhD<sup>3</sup>, Keith P. George PhD<sup>3</sup>, Jonathan Myers, PhD, Euan Ashley, MRCP DPhil<sup>1,2\*</sup>, François Haddad, MD<sup>1,2\*</sup>

1. Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, CA

2. Stanford Cardiovascular Institute, Stanford, CA

3. Liverpool, John Moores University, United Kingdom

\* Both authors are equivalent co-senior author of the study

**Key words:** Athletics, Level of Activity, Ventricular Remodeling and Function, Lean body mass, Deformation imaging,

**Address of correspondence:** Please address correspondence to Francois Haddad, MD FAHA, Stanford Cardiovascular Institute, Palo Alto, CA, 94304. Email: [fhaddad@stanford.edu](mailto:fhaddad@stanford.edu).

**Word Count: text (2927 words)**

## ABSTRACT

**Background:** There is limited data regarding ventricular remodeling in college female athletes, especially when appropriate scaling of cardiac dimensions to lean body mass (LBM) is considered. Moreover, it is not well established whether cardiac remodeling in female athletes is a balanced process with proportional increase in left ventricular (LV) mass and volume or right and left ventricular size.

**Methods:** During the pre-participation competitive screening, seventy-two college female athletes volunteered to undergo Dual Energy X-Ray Absorptiometry (DEXA) scan for quantification of LBM and comprehensive 2D echocardiography including assessment of longitudinal myocardial strain. The athletes were divided into two groups according to the intensity of the dynamic and static components of their sport categories, i.e. a higher intensity dynamic and resistive group (n=37 participating in rowing, water polo and La Crosse) and a lower intensity group (n=35, participating in short distance running, sailing, synchronized swimming and softball). In addition, we recruited a group of 31 age-matched non-athlete controls.

**Results:** The mean age of the study population was  $18.7 \pm 1.0$  years. When scaled to body surface area, the higher intensity group had  $17.1 \pm 3.6\%$  ( $p < 0.001$ ) greater LV mass when compared to the lower intensity group and  $21.7 \pm 4.0\%$  ( $P < 0.001$ ) greater LV mass than the control group. The differences persisted after scaling to LBM with  $14.2 \pm 3.2\%$  ( $p < 0.001$ ) greater LV mass in the higher intensity group. In contrast, there was no difference in any of the relative remodeling indices including the LV mass to volume ratio, right to left ventricular area ratio or left atrial to LV volume ratio ( $p > 0.50$  for all). In addition, no significant difference was noted among the three groups in LVEF ( $p = 0.22$ ), LV global longitudinal strain ( $p = 0.55$ ), LV systolic strain rate ( $p = 0.62$ ) or RV global longitudinal strain ( $p = 0.61$ ).

**Conclusion:** Female collegiate athletes participating in higher intensity dynamic and resistive sports have higher indexed LV mass even when scaled to LBM. The remodeling process does however appear to be a balanced process not only at the intraventricular level but also at the interventricular and atrio-ventricular levels.

## Introduction

Pre-participation screening of competitive college athletes is adopted by the majority of College programs and usually involves a medical history and physical examination, an electrocardiogram and in selected cases a focused echocardiogram. The interpretation of echocardiograms in athletes may however be challenging as the pattern of cardiac remodeling in athletes may vary according to sport type as well as lean body mass.

Morganroth et al.<sup>1</sup> were the first to suggest that athletic remodeling varies according to the intensity of the dynamic or static components of the sport discipline. In their landmark study, Morganroth et al. observed that male endurance trained athletes (with higher dynamic component) had a greater left ventricular (LV) end-diastolic dimension with a normal wall thickness (eccentric remodeling) whereas wrestlers (with higher static component) had increased wall thickness but normal LV end-diastolic dimension, hence more concentric remodeling. Recently, the “Morganroth” observation was challenged by recent studies suggesting that the increase in LV mass is proportional to the increase in LV volume (proportional remodeling) irrespective of the sport discipline.<sup>1-7</sup> Furthermore, difference in ventricular mass among sport discipline may be partially explained by differences in lean body mass (LBM) and not sport discipline.<sup>8,9</sup>

In contrast to the many studies focusing on cardiac remodeling in male college athletes, few studies have focused on female college athletes especially when appropriate scaling of cardiac dimensions to lean body mass (LBM) is considered.<sup>8,9</sup> For this study, we first hypothesized that college female athletes participating in higher intensity dynamic sports would have greater LV mass than lower intensity disciplines even after scaling to LBM. We further hypothesize that the cardiac remodeling in female athletes would be balanced irrespective of the intensity (dynamic) of the sport discipline, i.e. (1) LV mass would increase proportionally to ventricular volume, (2) right ventricular size proportionally to LV size, (3) atrial size proportionally to ventricular size and finally (4) right atria size proportionally left atria size. Finally, we

hypothesize that ventricular function including longitudinal strain would be comparable among sport disciplines irrespective of sport intensity.

## **METHODS**

As part of Stanford University athletic screening program, 75 college female athletes underwent pre-season cardiac screening using the American Heart Association (AHA)-12-point assessment, an electrocardiogram (ECG), a focused echocardiogram and a Dual Energy X-Ray Absorptiometry (DEXA)-based assessment of body composition. We excluded three African American athletes from analysis, as race is known to influence ventricular remodeling independent of sport category; none of athletes had an abnormal questionnaire, abnormal ECG or pathological echocardiographic findings leading to exclusion.<sup>10, 11</sup> We were able to divide participants into 2 groups according to dynamic or static component of their sport activity according to the classification of Mitchell et al.<sup>12</sup> (Figure 1), The first group included 37 subjects participating in higher intensity dynamic and static sport discipline including rowing, La Crosse or Water Polo (gray areas) while the second group included lower intensity dynamic sports including synchronized swimming, short distance track, softball, fencing and sailing. Although Water Polo is a sport discipline with moderate dynamic component, we chose to include it in the higher intensity group due to its higher static component and previous studies showing significant cardiac remodeling.<sup>13, 14</sup>

To further study the influence of physical activity on cardiac remodeling and function, we also recruited 32 healthy volunteers; one subject was subsequently excluded for mitral valve prolapse. The healthy volunteers had a normal AHA-12 points assessment, no family history of early cardiovascular disease and had to exercise less than 2 hours a week. Healthy volunteers did not, however undergo DEXA scans as this was only offered to athletes of the program.

### **Anthropometric measurements**

Lean body mass (LBM) was estimated using DEXA scan (Norland XR 26 Mark II/HS, Norland Corporation, WIS). Body surface area (BSA) was calculated with Dubois' formula. Body mass index

(BMI) was calculated using standard formula weight divided height squared (kg/m<sup>2</sup>).

### **Standard echocardiographic measurements**

All patients underwent standard transthoracic two-dimensional (2D) and color Doppler echocardiography using the Philips Sonos 7500 system (Philips Medical Imaging, Eindhoven, the Netherlands) and a 3.5-MHz transducer. The echocardiograms were blindly interpreted by an experienced reader (GG) according to the guidelines of the ASE.<sup>15</sup> LV mass was calculated in diastole using estimating LV mass based on the area-length (*AL*) formula (Figure 2).<sup>16</sup> LV end-diastolic and end-systolic volume (LVEDV and LVESV) were calculated using the 5/6 area length method to be consistent with the geometrical assumptions of the LV mass formula.<sup>16</sup> For quality control for measures of ventricular mass, we also measured ventricular mass in systole to verify that mass estimates were within 10% of each other; if not within 10%, a second reader (FH) read the studies and an average value of 2 readers was chosen (Figure 2). Atrial volumes were calculated using the apical 4 chamber views by the area-length method.<sup>16</sup>

Relative remodeling indices included: (1) LV mass to volume ratio (calculated using the 5/6 area length method), (2) relative LV wall thickness calculated using the average wall thickness and average ventricular diameter from the short axis view, (3) RV to LV end-diastolic area ratio and (4) relative atrial remodeling indices including LA volume to LV end-diastolic volume, RA to RV end-diastolic relative area and RA to LA relative area.

Measures of ventricular function included LV ejection fraction (LVEF) derived using the Simpson method in a 4 chamber view; stroke volume using the difference between end-diastolic and end-systolic volumes using the area-length methods; right ventricular (RV) fractional area change (RVFAC) and tricuspid annular plane systolic excursion (TAPSE) using 2D methodology. Analysis of LV and RV longitudinal strain (LS) were performed from the apical 4-chamber (4C) views, using vendor independent software (TOMTEC Imaging System, Unterschleissheim, Germany) as shown in Figure 2.<sup>17</sup> For LVLS, the 6 segments in the apical 4 chamber view were averaged while the 3 lateral segments were averaged for

RVLS measures. For the LV, strain rate parameters were also computed using the TOMTEC system. For ventricular strain measurements, the peak of the QRS was used as the reference point.

### **Scaling of cardiac dimensions**

Scaling of cardiac dimensions or mass was based on BSA and LBM. For BSA, the allometric coefficient chosen was unity based on the recommendations of the American Society of Echocardiography<sup>15</sup>; we chose to scale wall thickness and ventricular dimension to height as this is more consistent with the dimension of the parameter being scaled and likely more body size independent. For LBM, we chose the ones suggested by the studies of George et al. and Bella et al.<sup>8, 9, 18</sup> For LV mass and volume scaling to LBM was done using unity as the allometric coefficient, linear dimensions to LBM with an exponent of 0.33 and areas with an exponent of 0.66.<sup>8, 9, 18</sup>

### **Statistical analysis**

Results are expressed as mean  $\pm$  SD for normally distributed continuous variables or as number of cases and percentage for categorical variables. Comparison of groups was performed using Student's t-test for continuous variables assuming equal or unequal variance as appropriate and Chi-square test or Fisher Exact Test, as appropriate for categorical variables. The inter-observer variability was tested using the absolute and relative difference as well as the intraclass coefficient and coefficient of variation. Statistical analysis was performed using the PASW software (PASW 22.0 Inc, Chicago, IL).

## **RESULTS**

### **Population**

The demographic and anthropomorphic characteristics of the study population are summarized in table 1. There was no significant difference in age, BSA or BMI between the 3 groups. The percentage body fat were not statistically different between the higher intensity and lower intensity group. Athletes in the higher intensity group were, however taller and had a slightly higher values of LBM.

## **Differences in cardiac dimensions**

As shown in table 2, the HIGH group had significantly greater LV mass, LV volume and atrial volume when scaled to BSA. Even after scaling to LBM (table 3), there was still a significant difference in cardiac dimensions between the HIGH and LOW group. Expressed as relative differences, the LV mass was  $21.7 \pm 4.1\%$  and  $17.1 \pm 3.6\%$  greater in the HIGH compared to the LOW and CON (Figure 3a). In contrast there was no significant difference between groups in any of the relative remodeling indices including mass to volume ratio (relative difference of  $0.0 \pm 2.7\%$  and  $3.0 \pm 2.7\%$  and compared LOW and HIGH respectively), average relative wall thickness, RV to LV area, LAV to LVEDV or RA to LA relative area.

When using as a threshold the 95<sup>th</sup> percentile of the LOW group (reference group), there was a greater percentage of elevated LV mass when scaled to BSA, compared to LBM or mass to volume ratio (Figure 3b). The low percentage of abnormal value using the mass to volume ratio could be explained by the strong linear relationship between the LV mass and LV volume ( $R^2 = 0.71$ ,  $P < 0.001$ ) across sports type. Similar findings were found with regards to relative differences in LAV (Figure 3d, e and f).

With regards to the specific sport categories in each group, no significant difference in scaled ventricular mass was observed between rowing ( $2.46 \pm 0.36$  g/kg (rowing),  $2.54 \pm 0.36$  g/kg (water polo),  $2.39 \pm 0.22$  (lacrosse),  $P = 0.62$ ) in the HIGH group; no other difference in indexed ventricular mass was noted in the LOW groups.

## **Assessment of body size independent scaling parameters**

In our study, allometric scaling to LBM was associated with body size independent scaling in the HIGH and LOW groups. There was no significant relationship the LBM scaled dimension and LBM for LV mass:  $R^2 = 0.008$ ,  $P = 0.60$  (LOW) and  $R^2 = 0.05$ ,  $P = 0.14$  (HIGH); LVEDV,  $R^2 = 0.08$ ,  $P = 0.15$  (LOW) and  $R^2 = 0.006$ ,  $P = 0.65$  (HIGH) and LAV,  $R^2 = 0.003$ ,  $P = 0.75$  (LOW) and  $R^2 = 0.06$ ,  $P = 0.13$  (HIGH). In contrast, BSA did not yield body size independent scaling for LV mass in the high intensity athletes:  $R^2 = 0.14$ ,  $P = 0.02$



for LV mass,  $R^2 = 0.14$ ,  $P=0.02$  for LVEDV or and  $R^2 = 0.16$ ,  $P=0.01$  for LAV.

### **Ventricular function and Sport discipline**

Indexed stroke volume to BSA was  $17.6 \pm 3.6\%$  and  $19.6 \pm 4.4\%$  and higher in the HIGH when compared to the LOW and CON groups respectively and  $15.0 \pm 3.6\%$  higher in the HIGH group compared to the LOW group when scaled to LBM. There was no other significant difference in functional parameters in the HIGH group compared to the other groups (Table 5).

### **Inter-observer variability and quality control**

Inter-observer variability for LV mass measurements were measured using 15 athletes randomly selected (GG and YK). The absolute difference for LV mass was  $-6.4$  g  $[-17.1, 4.4]$  and the relative bias was  $-4.5\%$   $[-11.6, 2.6\%]$  with only one patient having a greater than 10% difference. The intraclass coefficient was 0.95 and the coefficient of variation of 3.7%. For quality control, we compared values of ventricular mass between diastole and systole. The mean difference between measures in systole and diastole was  $-3.1 \pm 5.4\%$  with only 12% of samples having differences more than 10% requiring averaging.

### **Discussion**

Our study has two main findings. First, it confirms that difference in LV mass in female athletes participating in high intensity sports persist even after scaling to lean body mass. Second, in contrast to the original Morganroth observation in male athletes, our study supports the balanced nature of ventricular remodeling in female athletes with proportional increase in ventricular mass and volume irrespective of sport type. In addition, we observed that the remodeling process is also balanced at the interventricular, atrio-ventricular and inter-atrial level.

Few studies have focused on ventricular remodeling in female athletes the majority being cross-sectional studies. In one of the largest echocardiography based studies, Caselli et al. analyzed ventricular remodeling in 148 female Olympic athletes using 3D echocardiography and scaling to BSA.<sup>2</sup> They found that endurance

(dynamic) and mixed (dynamic and static) sports had approximately 19% greater ventricular mass, and volume when compared to skilled (lower dynamic intensity) based sport categories.<sup>2</sup> Dressler et al. found slightly lower percentage increase in ventricular mass in the range of 11% in 276 female elite athletes<sup>19</sup> Five cross-sectional magnetic resonance (MRI) studies have compared ventricular remodeling in endurance female athletes compared to controls (n ranging from 20 to 59 athletes).<sup>5, 20-23</sup> These studies reported an increase in LV mass in the ranges from 19 to 26% when compared to on-athlete controls.<sup>5, 20-23</sup> Longitudinally, Howden et al. evaluated 5 female subjects who underwent endurance training, the change in ventricular remodeling that occur over 1 year in previously less active individuals and found that LV mass increased on average by 14%. Studies in male endurance using MRI find similar relative increase in LV mass. Prakken et al. and Luijkx et al. found that athletes participating in more dynamic sport activity have a 20 to 32% higher LV mass respectively.<sup>20, 22</sup> Echocardiography based studies in male athletes usually report greater differences in ventricular mass in the order of 39% likely due to an overestimation of wall thickness by echocardiography.<sup>24, 25</sup>

Our study represents the larger study scaling to LBM in college female athletes; as expected scaling to LBM reduces but does not eliminate the differences in LV mass or volume. Riley-Hagan et al. have previously shown in a smaller group of older female athletes participating in long distance running, cycling or cross-country skiing that endurance based disciplines had on average 17 to 19 % greater mass when compared to non-athletic controls.<sup>23</sup> The smaller difference of 14% observed in our study may be, in part, related to the pre-season timing of screening and by the fact that the athletes were not in the elite category.

One of the most important contribution of our study is the emphasis placed on the balance nature of cardiac remodeling in athletics. This challenges in part the Morganroth study observation that proposed that endurance based sports were associated with more eccentric remodeling.<sup>26</sup> In the study of Caselli et al. in Olympic based athletics, the mass to volume ratio appeared to be constant around one irrespective of the sport discipline.<sup>2</sup> These observation of balanced increase in LV mass and volume increase in athletes were confirmed by several cross-sectional and longitudinal based studies especially in male athletes.<sup>1, 20, 22,</sup>

<sup>27, 28</sup> Our study also highlights the balanced nature of the interventricular, atrio-ventricular and interatrial component of balanced remodeling. The balanced nature of ventricular remodeling does not appear to be limited to athletics. In fact, results from the Multi-Ethnic Study of Atherosclerosis (MESA) have previously shown that the mass to volume ratio is not influenced by level of activity although it is higher in older individuals, people of black race and in male subjects. As a reference, female subjects between 45 and 54 years in the MESA study had an average mass to volume ratio was 1.04.<sup>6</sup>

Functionally, sports with higher dynamic components have on average higher indexed stroke consistent with the higher cardiac output requirements often imposed by more aerobic-based physical activity. Consistent with the prior studies of Spence et al. and Andrea et al, no significant differences in LVEF or myocardial deformation indices were noted across sport discipline although the study was likely underpowered to show small differences.<sup>1, 29</sup> Moreover, and original to our study, no difference was observed for strain rate metrics among the different sport categories and controls.

Although not a major objective of our study, our study also highlights important methodological heterogeneity between the studies on ventricular remodeling. In fact, although the relative changes in mass or volume are consistent between the studies, the absolute values reported vary significantly. In our study, the measured values for both ventricular mass and mass to volume ratios in controls are comparable to the suggested reference cardiac MRI parameters of the study Kawel-Boehm et al.<sup>30</sup> but lower than previously reported echocardiographic studies.<sup>25</sup> In our study, measuring mass in both diastole and systole was associated with an excellent inter-reader reproducibility.

The clinical implications of the study are threefold. First, any future reference values for athletic screening should take into account sport category during screening. Second, the balanced nature of ventricular remodeling can be very useful when assessing for pathological remodeling in athletes. Building on this observation, Petersen et al. demonstrated that a maximal wall thickness to volume ratio of  $< 0.15 \text{ mm}/(\text{mL}/\text{m}^2)$  had a sensitivity of 80% and a specificity of 99% in differentiating pathological remodeling in

athletics and pathological hypertrophy with an AUC of 0.99.<sup>31</sup> In their study however, athletes were younger than patients with LV hypertrophy which may have led stronger discrimination.<sup>31</sup> Finally, future studies, validating the best methods for reproducibility between laboratories will be needed prior to recommending reference value for athletic screening.

The main limitation of our study is its small sample size and different sports type. However, despite its small size, the current study represents one of the largest among college female athletes; we did not also observe differences related to sport type. Although the sport classification could lead itself to some criticism, we based our categorization on the Task Force recommendation of Mitchell et al. and previous studies in athletes participating Water-Polo.<sup>12-14</sup> Another limitation is that we did not use magnetic resonance imaging for quantification of cardiac structure. However; our measures were performed by Stanford Cardiovascular Institute Biomarker and Phenotypic Core Laboratory by experienced readers with results are consistent with magnetic resonance based studies. Finally, we did not have complete training history and maximal oxygen consumption of participants. In addition, patients were not screened using the The Physical Activity Readiness Questionnaire for Everyone or the AHA/ACSM Health/Fitness Facility Pre-participation Screening Questionnaire.

In conclusion, these data highlight sport specific differences in ventricular and atrial remodeling in female college athletes. Mass to volume ratio or relative wall thickness is emerging as a parameter independent of sport discipline and will likely emerge as very useful for screening pathological remodeling.

## References

1. Spence AL, Naylor LH, Carter HH, Buck CL, Dembo L, Murray CP, Watson P, Oxborough D, George KP and Green DJ. A prospective randomised longitudinal MRI study of left ventricular adaptation to endurance and resistance exercise training in humans. *The Journal of physiology*. 2011;589:5443-52.
2. Caselli S, Di Paolo FM, Picicchio C, Di Pietro R, Quattrini FM, Di Giacinto B, Culasso F and Pelliccia A. Three-dimensional echocardiographic characterization of left ventricular remodeling in Olympic athletes. *The American journal of cardiology*. 2011;108:141-7.
3. Haykowsky MJ. Left ventricular remodelling and the athlete's heart: time to revisit the Morganroth hypothesis. *The Journal of physiology*. 2011;589:5915.
4. Naylor LH, George K, O'Driscoll G and Green DJ. The athlete's heart: a contemporary appraisal of the 'Morganroth hypothesis'. *Sports medicine*. 2008;38:69-90.
5. Steding K, Engblom H, Buhre T, Carlsson M, Mosen H, Wohlfart B and Arheden H. Relation between cardiac dimensions and peak oxygen uptake. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2010;12:8.
6. Turkbey EB, Jorgensen NW, Johnson WC, Bertoni AG, Polak JF, Diez Roux AV, Tracy RP, Lima JA and Bluemke DA. Physical activity and physiological cardiac remodelling in a community setting: the Multi-Ethnic Study of Atherosclerosis (MESA). *Heart*. 2010;96:42-8.
7. Lewis EJ, McKillop A and Banks L. The Morganroth hypothesis revisited: endurance exercise elicits eccentric hypertrophy of the heart. *The Journal of physiology*. 2012;590:2833-4.
8. George K, Sharma S, Batterham A, Whyte G and McKenna W. Allometric analysis of the association between cardiac dimensions and body size variables in 464 junior athletes. *Clinical science*. 2001;100:47-54.
9. George KP, Birch KM, Pennell DJ and Myerson SG. Magnetic-resonance-imaging-derived indices for the normalization of left ventricular morphology by body size. *Magnetic resonance imaging*. 2009;27:207-13.
10. Haddad F, Peter S, Hulme O, Liang D, Schnittger I, Puryear J, Gomari FA, Finocchiaro G, Myers J, Froelicher V, Garza D and Ashley EA. Race differences in ventricular remodeling and function among college football players. *The American journal of cardiology*. 2013;112:128-34.
11. Weinstock J and Estes NA, 3rd. The heart of an athlete: black, white, and shades of grey with no gold standard. *Circulation*. 2013;127:1757-9.
12. Mitchell JH, Haskell W, Snell P and Van Camp SP. Task Force 8: classification of sports. *Journal of the American College of Cardiology*. 2005;45:1364-7.
13. Pavlik G, Kemeny D, Kneffel Z, Petrekanits M, Horvath P and Sido Z. Echocardiographic data in hungarian top-level water polo players. *Medicine and science in sports and exercise*. 2005;37:323-8.
14. Horvath P, Petrekanits M, Gyore I, Kneffel Z, Varga-Pinter B and Pavlik G. Echocardiographic and spiroergometric data of elite Hungarian female water polo players. *Acta physiologica Hungarica*. 2009;96:449-57.
15. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, Flachskampf FA, Foster E, Goldstein SA, Kuznetsova T, Lancellotti P, Muraru D, Picard MH, Rietzschel ER, Rudski L, Spencer KT, Tsang W and Voigt JU. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *European heart journal cardiovascular Imaging*. 2015;16:233-70.
16. Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ, Chamber Quantification Writing G, American Society of Echocardiography's G, Standards C and European Association of E. Recommendations for chamber quantification: a report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing

- Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *Journal of the American Society of Echocardiography : official publication of the American Society of Echocardiography*. 2005;18:1440-63.
17. Voigt JU, Pedrizzetti G, Lysyansky P, Marwick TH, Houle H, Baumann R, Pedri S, Ito Y, Abe Y, Metz S, Song JH, Hamilton J, Sengupta PP, Kolias TJ, d'Hooge J, Aurigemma GP, Thomas JD and Badano LP. Definitions for a common standard for 2D speckle tracking echocardiography: consensus document of the EACVI/ASE/Industry Task Force to standardize deformation imaging. *European heart journal cardiovascular Imaging*. 2015;16:1-11.
  18. Bella JN, Devereux RB, Roman MJ, O'Grady MJ, Welty TK, Lee ET, Fabsitz RR and Howard BV. Relations of left ventricular mass to fat-free and adipose body mass: the strong heart study. The Strong Heart Study Investigators. *Circulation*. 1998;98:2538-44.
  19. Pressler A, Haller B, Scherr J, Heitkamp D, Esefeld K, Boscheri A, Wolfarth B and Halle M. Association of body composition and left ventricular dimensions in elite athletes. *European journal of preventive cardiology*. 2012;19:1194-204.
  20. Luijckx T, Cramer MJ, Prakken NH, Buckens CF, Mosterd A, Rienks R, Backx FJ, Mali WP and Velthuis BK. Sport category is an important determinant of cardiac adaptation: an MRI study. *British journal of sports medicine*. 2012;46:1119-24.
  21. Petersen SE, Hudsmith LE, Robson MD, Doll HA, Francis JM, Wiesmann F, Jung BA, Hennig J, Watkins H and Neubauer S. Sex-specific characteristics of cardiac function, geometry, and mass in young adult elite athletes. *Journal of magnetic resonance imaging : JMRI*. 2006;24:297-303.
  22. Prakken NH, Velthuis BK, Bosker AC, Mosterd A, Teske AJ, Mali WP and Cramer MJ. Relationship of ventricular and atrial dilatation to valvular function in endurance athletes. *British journal of sports medicine*. 2011;45:178-84.
  23. Riley-Hagan M, Peshock RM, Stray-Gundersen J, Katz J, Ryschon TW and Mitchell JH. Left ventricular dimensions and mass using magnetic resonance imaging in female endurance athletes. *The American journal of cardiology*. 1992;69:1067-74.
  24. Prakken NH, Teske AJ, Cramer MJ, Mosterd A, Bosker AC, Mali WP, Doevendans PA and Velthuis BK. Head-to-head comparison between echocardiography and cardiac MRI in the evaluation of the athlete's heart. *British journal of sports medicine*. 2012;46:348-54.
  25. Utomi V, Oxborough D, Whyte GP, Somauroo J, Sharma S, Shave R, Atkinson G and George K. Systematic review and meta-analysis of training mode, imaging modality and body size influences on the morphology and function of the male athlete's heart. *Heart*. 2013;99:1727-33.
  26. Morganroth J, Maron BJ, Henry WL and Epstein SE. Comparative left ventricular dimensions in trained athletes. *Annals of internal medicine*. 1975;82:521-4.
  27. Scharhag J, Schneider G, Urhausen A, Rochette V, Kramann B and Kindermann W. Athlete's heart: right and left ventricular mass and function in male endurance athletes and untrained individuals determined by magnetic resonance imaging. *Journal of the American College of Cardiology*. 2002;40:1856-63.
  28. Spence AL, Carter HH, Murray CP, Oxborough D, Naylor LH, George KP and Green DJ. Magnetic resonance imaging-derived right ventricular adaptations to endurance versus resistance training. *Medicine and science in sports and exercise*. 2013;45:534-41.
  29. D'Andrea A, Cocchia R, Riegler L, Scarafilo R, Salerno G, Gravino R, Golia E, Pezzullo E, Citro R, Limongelli G, Pacileo G, Cuomo S, Caso P, Russo MG, Bossone E and Calabro R. Left ventricular myocardial velocities and deformation indexes in top-level athletes. *Journal of the American Society of Echocardiography : official publication of the American Society of Echocardiography*. 2010;23:1281-8.
  30. Kawel-Boehm N, Maceira A, Valsangiacomo-Buechel ER, Vogel-Claussen J, Turkbey EB, Williams R, Plein S, Tee M, Eng J and Bluemke DA. Normal values for cardiovascular magnetic resonance in adults

and children. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2015;17:29.

31. Petersen SE, Selvanayagam JB, Francis JM, Myerson SG, Wiesmann F, Robson MD, Ostman-Smith I, Casadei B, Watkins H and Neubauer S. Differentiation of athlete's heart from pathological forms of cardiac hypertrophy by means of geometric indices derived from cardiovascular magnetic resonance. *Journal of cardiovascular magnetic resonance : official journal of the Society for Cardiovascular Magnetic Resonance*. 2005;7:551-8.

## Figure legends

**Figure 1.** Sport discipline classification based on oxygen uptake and maximal voluntary contraction (MVC). The gray shaded squares represent the group participating in higher dynamic-static component sports (HIGH) versus the lower dynamic-static component sports.

**Figure 2. Representative measures for LV remodeling and myocardial deformation analysis.** Panel A through C highlight the different components for measuring LV mass using the area length method; in our study, we measured mass in both diastole and systole integrating this for quality control method (conservation of mass principle). Panels D through F highlight the different longitudinal deformation metrics used in the study; E shows the average myocardial longitudinal strain curve and F the strain rate metrics. LS indicates longitudinal strain; SR, strain rate.

**Figure 3. Relative difference in cardiac remodeling indices in the athletic and control groups.** Panel A shows the difference in LV mass when scaled to BSA and LBM in the higher intensity sport group (HIGH) group compared to the control group (C) and the Lower intensity sport group (L) as well as for the mass to volume ratio (M/V). Panel B shows percentage of athletes presenting a value higher than the 95% of value in the lower intensity group; in our study the limit chosen for LVM indexed to BSA is consistent with the upper limit in the recent MRI study of Kawel-Boehm.<sup>30</sup> Panel C shows the strong linear relationship between LV mass and volume in the 3 groups. Panel D, E and F shows corresponding analysis for atrial remodeling.