

**Leonardi, N, Carnacina, I, Donatelli, C, Ganju, NK, Plater, AJ, Schuerch, M and Temmerman, S**

**Dynamic interactions between coastal storms and salt marshes: A review**

**<http://researchonline.ljmu.ac.uk/id/eprint/9345/>**

#### **Article**

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Leonardi, N, Carnacina, I, Donatelli, C, Ganju, NK, Plater, AJ, Schuerch, M and Temmerman, S (2017) Dynamic interactions between coastal storms and salt marshes: A review. *Geomorphology*, 301 (15). pp. 92-107. ISSN 0169-555X**

LJMU has developed **[LJMU Research Online](#)** for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

# **Dynamic interactions between coastal storms and salt marshes: a review**

Nicoletta Leonardi (a\*), Iacopo Carnacina (b), Carmine Donatelli (a), Neil Kamal Ganju (c),  
Andrew James Plater (a), Mark Schuerch (d), Stijn Temmerman (e)

\* Corresponding author nicleona@liverpool.ac.uk

(a) University of Liverpool, Department of Geography and Planning, 74 Bedford St S.  
Liverpool, UK

(b) Liverpool John Moores University, Department of Civil Engineering, Peter Jost Enterprise  
Centre, Byrom Street, Liverpool, L3 3AF

(c) U.S. Geological Survey, Woods Hole Coastal and Marine Science Center, MA 02543, USA

(d) Cambridge Coastal Research Unit (CCRU) Department of Geography, University of  
Cambridge, Downing Place, Cambridge, CB2 3EN, United Kingdom

(e) Ecosystem Management Research Group, University of Antwerp, Universiteitsplein 1,  
2610 Wilrijk, Belgium

## **Abstract**

The action of storms, and associated large waves and inundation depths, can strongly  
alter horizontal and vertical salt marsh dynamics in the immediate after-storm period, as well  
as in the longer term. This manuscript reviews the progresses made in the understanding of the  
dynamic interactions between coastal storms and salt marshes, including the dissipation of  
extreme water levels and wind waves across marsh surfaces, the geomorphic impact of storms  
on salt marshes, the preservation of hurricanes signals and deposits into the sedimentary  
records, and the importance of storms for the long term survival of salt marshes to sea level  
rise. Salt marshes are effective in dissipating wave energy, and storm surges, especially when

the marsh is highly elevated, continuous, and more than 10km wide. This buffering action, is very effective during moderate storms, but less efficient for long storms lasting more than one day; for this reason the use of hybrid approaches, combining continuous marshes with engineered defence structures is recommended for coastal protection. From a morphological point of view, our considerations highlight the necessity to focus on the indirect long term impact that large storms exerts on the whole marsh complex rather than on sole after-storm periods. Storms can cause tidal flats deepening which in turn promotes wave energy propagation, and exerts a long term detrimental effect for marsh boundaries even during calm weather. On the other hand, when a violent storm causes substantial erosion but sediments are redistributed across nearby areas, the long term impact might not be as severe as if sediments were permanently lost from the system, and the salt marsh could easily recover.

## **1. Introduction**

### **1.1 Changing storm activity**

Many areas are experiencing a change in both extreme and mean storm conditions as a consequence of a changing climate (e.g. Zhang et al., 2000; Webster et al., 2005; Bacmeister et al., 2016). For example, according to the Intergovernmental Panel on Climate Change (IPCC, e.g. Solom et al., 2007; Pachauri et al., 2014) it is virtually certain (99-100% probability) that the intensity of cyclone activity has increased in the North Atlantic since 1970, even if there is low confidence that the long term changes are robust. In terms of extremes, it is likely (66-100% probability) that extreme sea levels such as the ones experienced during storm surges have increased since 1970 on a global average. The latter trend has been mainly attributed to an increase in mean sea level even if more studies are necessary to fully separate the effect of global mean sea level rise from the effects of more local modifications to the coastal systems

(e.g. Pachauri et al., 2014). Finally, it is also likely that there are more land regions where the number of heavy precipitation events has increased than where it has decreased.

Evaluations of future increases in storms and hurricanes activity are complex, and with large uncertainties. For example, a statistical correlation has been found between the power dissipation index of hurricanes (i.e. an index combining intensity, frequency and duration of hurricanes) and Atlantic Sea Surface Temperature (SST) (e.g. Vecchi et al., 2008). Based on this relationship and taking into account hurricanes activity since 1950, as well as future SST projection, there should be a 300% increase in hurricanes activity by the late 21<sup>st</sup> century. However, a statistical correlation has been also found between the power dissipation index and the Atlantic sea surface temperature relative to the Tropical mean sea temperature; if the latter relationship is considered, the projected change in hurricane activity by 2100 would be around 25%, which is modest with respect to the estimation above (Vecchi et al., 2008). Projections about the future of hurricanes activity might get even more complicated when looking at the longer term. Mean air temperature, Atlantic SST and the unadjusted hurricanes count all show a marked increase since the late 1800; however, when the raw hurricane count is adjusted for the storms which were not counted during the pre-satellite era due to technology, and ship track density limitations, no significant increase is observed (e.g. Vecchi et al., 2008).

Figure 1 illustrates model results in relation to the 21<sup>st</sup> century changes in Emmanuel's (1995) wind maximum potential intensity (MPI<sub>v</sub>), the increase of which is generally associated with an increase in storms activity and intensity (Vecchi and Sobel, 2007). Results refer to the IPCC-AR4 Scenario A1B for the period from June-November. The MPI<sub>v</sub> index increases over most of the northern hemisphere and tropical zone of the southern hemisphere, but there are also large areas particularly in the southern hemisphere indicating decreases. The regions where the MPI<sub>v</sub> decreases are associated with a relative minimum in SST (e.g. Sobel et al., 2002).

On a regional scale, for instance, by using a barotropic type surge model and global conditions representative of the IPCC A2 SRES scenarios between 1961-1990 and 2071-2100, it was shown that storm surge extremes may also significantly increase along most of the North Sea coast toward the end of this century (Woth et al., 2006). Recent results from ensemble simulation runs using Regional Climate Models for various locations in the United States (Jiang et al., 2016) also support the hypothesis of variations in future storm pattern; specifically, they predict shorter storm durations, longer inter-storms periods, and higher storms intensities.

In spite of the abundance of studies in relation to climatic projections and past trends, many challenges are still present, especially for the monitoring of coastal zones, due to limitations of some current modelling and field practice frameworks. For instance, the retrieval of waves and winds in the coastal areas is not yet as mature as sea level measurements, and the development of a wider applicability of altimetry techniques could be relevant for the simultaneous monitoring of wave height, wind speed and sea levels. In this context, Liu et al. (2012) showed the potential usefulness of the 1-Hz along-track altimetry data for the description of shelf areas, and Passaro et al., 2015 showed that estimations of wave height from ALES (Adaptive Leading Edge Sub-waveform retracker) were better correlated to buoy data than processed products. Such techniques could be coupled to standard modelling, and field data approach to build a more comprehensive and homogeneous database for the study of these coastal ecosystems

## **1.2. Pressures on salt marsh ecosystems**

Salt marshes are important coastal ecosystems frequently fringing the interior of estuaries and bays, and establishing in low-energy inter-tidal zones. Due to their location and vegetated surfaces, salt marshes offer several ecosystem services. For example, their value for buffering against the impact of storms has been estimated up to 5 million USD per km<sup>2</sup> in the United States (e.g., Costanza et al., 2008), and 786 million GBP per year for UK marshes (UK

National Ecosystem assessment, 2011; Foster et al., 2013; Moller et al., 2014). Indeed, in recent years, salt marsh conservation and restoration projects are increasingly adopted as part of coastal and estuarine flood defence programs, based on the concept of “living shorelines” or “nature-based solutions” for flood defence (e.g., Temmerman et al., 2013; Fagherazzi, 2014).

Apart from flood protection, other salt marsh services include the storage of sediments, pollutants, nutrients, as well as of large amounts of carbon at a geological time scale (e.g. Mudd, et al., 2009; Kirwan and Mudd, 2012; Pendleton et al., 2012). They are also the natural habitat of many plants and animal communities, and offer a place for recreational and touristic activities (e.g. Barbier et al., 2011).

The long-term persistence of salt marshes appears related to the maintenance of a delicate balance between sediment and nutrient inputs, and external agents such as wave energy, storm surges, tidal inundation, and sea level rise (e.g. Spencer et al., 1998; Plater et al., 1999; van de Koppel et al., 2005; Deegan et al., 2012; Fagherazzi et al., 2012; Kirwan et al., 2016; Leonardi et al., 2016). Figure 2 represents a sketch of some of the main physical and ecological processes acting on a salt marsh. This includes, for instance, the exchange of sediments between the tidal flat and the marsh platform, biomass production and sediment deposition on the marsh platform promoting vertical accretion, and possible erosion/progradation of the marsh edge. Ultimately, the survival of salt marshes has been related to a sediment budget problem (Ganju et al., 2017).

Salt marshes have been found to be extremely vulnerable, and large salt marsh losses have been documented worldwide. For instance, for areas in the south west of the Netherlands and the Wadden Sea, marsh edge erosion rates up to 4 m/yr have been observed, in spite of vertical accretion rates in balance with sea level rise (e.g., Bakker et al., 1993). In England and Wales salt marsh areal loss has been estimated to be around 83 ha yr<sup>-1</sup> (Environment Agency, 2011; Foster et al., 2013), 105 ha yr<sup>-1</sup> for the period in between 1993 and 2013 (Pye and French,

1993), and is projected to be 349 ha yr<sup>-1</sup> for the period between 1998 and 2048 (Lee, 2001). In the Greater Thames area, the erosion was estimated to be around 25% of the total area present in 1973 (Cooper et al., 2009), while in the Solent (UK) 40% of the total salt marsh area present in 1971 was eroded between 1971 and 2001 (Cope et al., 2008). Erosion up to 80 cm/yr has been recently measured in the northern part of the Venice Lagoon (e.g., Bondoni et al., 2016). For the East Coast of the United States, in Plum Sound and the Virginia Coast Reserve, salt marsh boundary erosion rates ranged from a couple of cm up to 3 m/yr over a 7-year measuring period (Leonardi and Fagherazzi, 2014, 2015). In Barnegat Bay, New Jersey, USA, erosion rates from 1930 to 2007, and from 2007 to 2013, were similar, with around half of the marsh area that fringes the interior of the bay eroding less than 0.5 m/yr, the other half displaying erosion rates up to 2 m/yr, and only a 3 percent eroding faster than 2 m/yr (Leonardi et al., 2016b). A recent global analysis on salt marsh erosion and wave measurements by Leonardi et al., 2016a revealed that most of salt marsh deterioration is caused by moderate storms of a monthly frequency while intense hurricanes contribute to less than 1% to long term salt marsh erosion rates.

The action of storms and associated wind waves and storm surges can strongly alter both horizontal and vertical salt marsh dynamics in the immediate after-storm period, as well as in the long term, by affecting erosion/ deposition, and sediment import/ export in salt marshes and surrounding areas. Furthermore, storms generate serious flood risks in low-lying and highly populated coastal zones. For these reasons, and especially under a climate change perspective, it is important to understand the reciprocal interaction between storms and salt marshes. This manuscript aims to review progresses made in the understanding of salt marsh-storms interactions, and is organized as follows: we first review storm surges (section 2), and wind waves (section 3) attenuation across salt marshes. In section 4 we focus on the impact of storms on salt marshes morphology, and on the preservation of hurricanes signals into the

sedimentary records. Section 5 focuses on the impact of storms on the marsh sediment budget. Section 7 discusses how the interplay between storms occurrence and sea level rise influences salt marsh survival. A set of discussions and conclusions is finally presented.

## **2. Storm surge attenuation by salt marsh**

Vegetated coastal ecosystems, in particular salt marshes and mangroves, are increasingly valued for their protective function against storm surge flood risks. This is illustrated by the rapidly increasing number of scientific studies on storm surge attenuation by salt marshes and mangroves, and growing societal interest in so-called ecosystem-based or nature-based flood defence programs, i.e. marsh and mangrove restoration projects aiming to mitigate storm surge flood risks (e.g. Cheong et al., 2013; Sutton-Grier et al., 2015; Temmerman et al., 2013). The effectiveness of storm surge height reduction behind marshes is commonly quantified as the attenuation rate in cm of surge height reduction per km distance that the storm surge has propagated over marshes (e.g. Wamsley et al., 2010). However, mechanistic insights in the various factors that control this attenuation rate are rather fragmentary presented in recent literature, which may be one reason why real life implementations of nature-based flood defence are relatively scarce so far (Temmerman et al., 2013). Here in this section, we review the most recent scientific insights.

Although anecdotal evidence of storm surge protection behind large marshes is presented in early reports (e.g. Lovelace, 1994; USACE, 1963), systematic evidence and mechanistic studies only started to accumulate over the past 10 years. In particular major coastal flood disasters caused by the Indian Ocean tsunami in 2004 and hurricane Katrina along the US Gulf coast in 2005 boosted worldwide scientific and public awareness of the potentially important protective role of mangroves (Danielsen et al., 2005) and salt marshes (Day et al., 2007). A first important source of empirical evidence comes from studies that analysed the



reduction of damage or human deaths as a function of marsh or mangrove width between coastal settlements and the open sea. For example, Costanza et al., 2008, performed an extensive analysis of 34 major hurricanes that hit the US Atlantic and Gulf coasts since 1980, demonstrating that damage to properties was significantly reduced behind marshes, and that a loss of 1 ha of marshes would increase average storm damages by 33000 USD. For mangroves, Das and Vincent, 2009, showed that villages that were hit by a tropical cyclone surge in India experienced significantly lower numbers of deaths when they had wider mangroves between them and the coast.

A second source of empirical evidence, are direct measurements of storm surge height reduction within and behind large marshes. Data reported in the literature are especially from the US Gulf coast (e.g. Lovelace, 1994; McGee et al., 2006; USACE, 1963), which is regularly hit by hurricane storm surges and where huge marshlands of several tens of kilometres wide exist in the Mississippi delta and in back-barrier tidal lagoons. A rule of thumb, derived from these reports, is that peak surge levels are reduced by on average 1 m for every 14.5 km that the surge has propagated over marshes (i.e.  $\sim 6.9$  cm/km), with large variations between individual hurricane events as much as from 1 m surge reduction per 4 km of marshland (i.e. 25 cm/km) to only 1 m per 60 km (i.e.  $\sim 1.7$  cm/km) (based on data compilation by Wamsley et al., 2010). This large variation in empirical data indicates that storm surge propagation and attenuation over marshes is complex and that the effectiveness of surge height reduction largely varies depending on specific storm characteristics, marsh ecosystem properties and larger-scale coastal landscape settings. For a macro-tidal estuarine marsh in the SW Netherlands, Stark et al., 2015, presented a large dataset ranging from regular tides to storm surges, showing that the magnitude of tidal and storm tide attenuation strongly depends on the marsh inundation depth and the dimensions of channels that dissect the marsh landscape. Maximum attenuation rates of up to 5 cm/km were measured over marsh transects with smaller channels and for marsh

inundation depths of 0.5-1 m, while attenuation rates decreased for shallower and deeper inundation events, including storm surges. For mangroves in Southern Florida, hurricane surge attenuation rates of 9.4 cm/km have been measured over relatively continuous mangrove forests, and slightly lower rates for mangroves along a river corridor (Krauss et al., 2009).

Hydrodynamic modelling studies are a third line of evidence and important research tools to disentangle the various factors controlling the effectiveness of storm surge height reduction by wetlands. Comparing the rapidly growing number of publications in the past few years (see below), we can generally make a distinction between two main mechanisms that depend on the larger-scale landscape setting: (1) storm surge attenuation within and behind continuous marshes is basically due to *friction* exerted by the marsh vegetation and soil on the landward propagating storm surge (e.g. Sheng et al., 2012); and (2) storm surges propagating through an estuarine or deltaic channel or embayment can be attenuated due to lateral flooding and *water storage* on marshes adjacent to that channel (e.g. Smolders et al., 2015). The frictional effect (1) is called here *within-marsh attenuation* and the water storage effect (2) *along-channel attenuation*. Ultimately both take place in most real cases, as marshes and mangroves are typically dissected by networks of tidal channels, implying that surge propagation along these channels is affected by both frictional and lateral water storage effects (e.g. Stark et al., 2016).

Modelling studies, either for idealized marsh geometries (e.g. Loder et al., 2009; Sheng et al., 2012; Temmerman et al., 2012) or for specific more realistic landscape settings (e.g. Resio and Westerink, 2008; Wamsley et al., 2010; Wamsley et al., 2009; Zhang et al., 2012), demonstrate that the effectiveness of storm surge attenuation depends on specific properties of (1) the storm forcing (such as storm intensity, duration, forward moving speed, storm track), (2) the marsh ecosystem (such as marsh size and soil elevation, vegetation density and continuity, within-marsh channel dimensions), and (3) larger-scale coastal landscape settings

(such as off-shore bathymetry, shoreline shape, open coast, back-barrier, estuarine or deltaic setting, levees or dikes behind marshes, etc.).

In terms of effects of storm characteristics, attenuation rates are generally higher for shallow to moderate storm surge levels and decrease for more extreme storm surges that deeply submerge the marshes, as within-marsh frictional effects on the storm surge attenuation relatively decrease with increasing water depth on the marsh (Lawler et al., 2016; Resio and Westerink, 2008; Sheng et al., 2012; Wamsley et al., 2010). Similarly, marshes with a higher soil elevation are more effective in attenuating higher storm surges (Loder et al., 2009; Smolders et al., 2015; Stark et al., 2016), implying that marshes with a sediment accretion deficit and consequently decreasing surface elevation relative to rising sea level, lose their effectiveness for storm surge protection (Temmerman et al., 2012; Wamsley et al., 2009). The protective function also decreases for storms with a longer duration, as the surge has more time to propagate landward and to fill up the whole marsh area (Resio and Westerink, 2008; Wamsley et al., 2010). Similarly, storm surge attenuation behind wetlands is more effective for storms with a faster forward moving speed (Hu et al., 2015; Liu et al., 2013; Sheng et al., 2012; Zhang et al., 2012).

In terms of marsh ecosystem properties, obviously wider marshes, of at least 10 or more kilometres wide, are more effective, as well as marshes with a higher soil elevation, as explained above. Effectiveness of storm surge attenuation also markedly increases when marsh vegetation is simulated that exerts more friction (Hu et al., 2015; Loder et al., 2009; Sheng et al., 2012), and with higher ratios of marsh vegetation to open water (Temmerman et al., 2012; Zhang et al., 2012), implying that patchy patterns of gradual marsh degradation, which are observed in several marshes around the world (e.g. Schepers et al., 2017), lead to loss of their storm protection function (Temmerman et al., 2012). The dimensions of channels, which typically cut into marshes, play a major role: simulations with deeper or wider channels, show

that landward flood propagation through the channels is facilitated leading to less storm surge height reduction (Stark et al., 2016; Temmerman et al., 2012). (Stark et al., 2016) showed for a marsh in the SW Netherlands that the effects of within-marsh channel dimensions, marsh platform elevation and storm surge height can be combined into one parameter predicting variations in attenuation rate from 0 to nearly 25 cm/km, i.e. as a function of the ratio between the water volume that is present at high tide above the marsh platform and the total water volume above the platform and in the channels (Figure 3).

Finally, the precise rates of storm surge attenuation by marshes depend on case-specific larger-scale landscape settings. For example, significant storm surge attenuation by wetlands is simulated for the several tens of kilometres wide marshes in the Mississippi deltaic area (Barbier et al., 2013; Hu et al., 2015; Resio and Westerink, 2008; Wamsley et al., 2010; Wamsley et al., 2009) and wide mangrove systems in Southern Florida (Liu et al., 2013; Zhang et al., 2012), while more moderate to limited contribution of marshes to storm surge protection are simulated for marshes along the Chesapeake Bay (Haddad et al., 2016), and back-barrier lagoon systems of Jamaica Bay, New York (Marsooli et al., 2016) and the Delmarva coast (Lawler et al., 2016). For the case of marshes occurring along the funnel shaped Scheldt estuary in the Netherlands and Belgium, simulations show that marshes of the same size but located more upstream are more effective in attenuating storm surges propagating upstream along the estuarine channel (Smolders et al., 2015). Man-made structures, in particular coastal defence structures such as levees and dikes behind marshes, may cause the setup of water levels against these structures and hence limit the storm surge attenuating effect of marshes in front of such structures, as shown for example in simulations for the 2005 hurricanes Katrina and Rita in the Mississippi delta (Wamsley et al., 2009). Similarly, for a marsh in the SW Netherlands, (Stark et al., 2016) showed blockage effects and setup of peak surge levels against dikes behind the

marsh, and that the marsh width needs to be at least 6-10 km to avoid such blockage effects and to maximize the rate of storm surge attenuation.

Summarizing, we may say that empirical data and modelling studies demonstrate effective storm surge height reduction behind large (at least 10 km wide), high-elevated and continuous marshes with few or small channels, and by marshes located more inland along funnel-shaped estuarine and deltaic channels, especially during moderate storm surges, but less effectively during extreme storms that continue for more than a day. The latter implies that solely relying on nature-based flood defence in populated low-lying coastal and estuarine areas is commonly not advised. Instead so-called hybrid approaches, combining conservation and restoration of continuous marshes with engineered defence structures, are increasingly developed and implemented worldwide (Sutton-Grier et al., 2015; Temmerman and Kirwan, 2015; Van Wesenbeeck et al., 2014), e.g. on large scales in the Mississippi delta (CPRA, 2012) and Scheldt estuary in Belgium (Meire et al., 2014). An important argument for such hybrid approaches, is that they are more cost-effective as they do not only provide flood risk mitigation but also other valuable ecosystem services, and marshes and mangroves build up land with rising sea levels, making them self-adaptive defences in face of global change (Temmerman et al., 2013).

### **3. Wave energy dissipation by salt marsh**

Salt marshes are natural wave energy dampers (e.g. Moeller, 2006; Moeller et al., 2014; Spencer et al., 2016). For shallow water, the dissipation of wave energy is related to the viscous boundary layer friction, permeability, and viscous layer of the seabed (e.g. Le Hir et al., 2000). Over a salt marsh the bed-roughness might be considered as the result of two contributions, i.e., vegetation induced friction, and topographic variations over the marsh surface (Hartnall, 1984; Dijkema, 1987; Pethick, 1992). It is also recognized that wave attenuation is affected by

plant characteristics such as geometry, stem density, spatial coverage, and stiffness, and that hydrodynamic conditions such as water depth, wave period, and wave height are relevant.

The pioneer work conducted in relation to the interaction between wave oscillatory motion and vegetation has been mainly aimed at quantifying wave attenuation within vegetation (e.g. Fonseca and Cahalan, 1992; Kobayashi et al., 1993). Standard approaches for the prediction of wave energy attenuation by vegetation, are based on the equation for the conservation of energy where the local flow field is estimated using linear wave theory. This approach, while reasonable, might be compromised if the vegetation substantially modifies the flow field. An alternative approach was proposed by Kobayashi et al., 1993, for the submerged vegetation case, for which the problem was formulated by using the continuity and linearized momentum equations for the regions over and within the vegetation canopy. By considering the effect of vegetation in terms of drag coefficient, introducing an unknown damping coefficient, and linearizing the friction term, they obtained an analytical solution for small monochromatic waves whose amplitude has been found to decay exponentially in the propagation direction. Koch and Gust, 1999, suggested that the periodic motion of seagrass blades also promotes mass transfer between the meadow, and the overlying water column. Luhar et al., 2010, demonstrated that even when the motion is driven by a purely oscillatory flow, a mean current in the direction of wave propagation is generated within the meadow. This current is forced by non-zero wave stress similar to the streaming observed in wave boundary layers, and the current is approximately four times the one predicted by the laminar boundary layer theory.

Among others, the dissipation of wind waves has been found to increase with increasing relative wave height, i.e. the ratio between wave height and water depth (e.g. Le Hir et al., 2000). Field measurements in England support this relationship, and show that for the analyzed field sites the relationship was mainly valid for relative wave height ratios above a critical

lower limit and below 0.55; when the ratio is below the lower limit, waves become too small (or water depth too high) to have an effective vegetation-wave interaction; however, when the relative wave height is  $> 0.55$ , the relationship between wave dissipation and relative wave height becomes invalid because the maximum dissipation capacity of vegetation has been reached (Moeller, 2006).

Field measurements of wind waves over sand flat to salt marsh cross-shore transects, also suggest that wave energy dissipation over salt marshes is significantly higher (up to 82% of the energy is dissipated) than on sand flats (29% dissipation) (Moeller, 1999, Figure 4). While part of the wave damping effect is attributable to the reduction in water depth on the higher elevated marsh platform (relative to the lower elevated tidal flat), the energy dissipation over salt marshes is up to 50 % stronger even under similar water depth conditions, which proves the important role of vegetation in the dissipation process.

Another parameter controlling the rate of energy dissipation is the ratio between water depth and plants height (submergence ratio, i.e. Yang et al., 2012): the smaller this ratio, the larger the wave attenuation rate (Augustin et al., 2009; Paul et al., 2012). Wave damping is also strictly related to the relative motion between fluid and plants, which depends on plants stems flexibility, stems diameter, and stems length. Stems with relatively high stiffness tend to follow an oscillatory swaying movement throughout the wave cycle, while more flexible stems tend to bend in the dominant direction of the orbital flow with a high angle which results in canopy flattening, and loss of flow resistance (whip-like movement) (i.e. Luhar and Nepf, 2016; Mullarney and Henderson, 2010; Paul et al., 2016). The movement can switch from swaying to whip-like as the wave energy increases (for example during storm periods) (e.g. Luhar and Nepf, 2016). Increasing plant flexibility reduces the damping of waves as stems tend to move with the surrounding water (Bouma et al., 2005; Elwany et al., 1995; Riffe et al., 2011), however stiff plants can break if hydrodynamic loads are higher than a critical value (Heuner

et al., 2015; Puijalon et al., 2011; Silinski et al., 2015). The dissipative contribution given by flexible plants is low, but their deformed configuration (flattening) under high orbital velocities ( $\geq 74 \text{ cm s}^{-1}$ ) helps to stabilize surface sediments (Neumeier and Ciavola, 2004; Peralta et al., 2008). In contrast, more rigid plants can reach breakage (from medium orbital velocities), increase turbulence and sediment scouring around the stems (reference) and cause more erosion due to increased shear stress values (Spencer et al., 2016).

During extreme storms and associated storm surges, waves and water levels are the highest, and hence it can be questioned whether, under these conditions, salt marshes still play a considerable role in wave attenuation. Large scale laboratory experiments (Moeller et al., 2014) confirm that, even under extreme conditions, wave energy dissipation by salt marshes is very high, and up to 60% of this wave energy reduction is attributed to the presence of vegetation. As the storm progresses, vegetation stems are gradually flattened and the wave dissipation decreases, but as suggested by previous work (e.g. Neumeier and Ciavola, 2004; Peralta et al., 2008), the flattening of vegetation promotes the stability of the substrate. Paul et al., 2016 tested different artificial vegetation elements to measure drag forces on vegetation under different wave loading. They found that stiffness and dynamic frontal areas (e.g. frontal area resulting from bending) are the main factors determining drag forces, while the still frontal area of plants dominate the force-velocity relationship only for low orbital velocities. Rupprecht et al., 2015 presented biophysical properties of species commonly found in NW European salt marshes, and compared the performance of two methods for the non-destructive assessment of aboveground biomass during storms, i.e. measurements of light availability within vegetation canopy, and side-on photography vegetation, with the latter being found more accurate. In the same experiments as reported by Moeller et al. 2014, Rupprecht et al., 2017, tested the effectiveness of two typical NW European salt marsh grasses (*Puccinellia maritima*, and *Elymus athericus*) under simulated storms and no-storms conditions. They found



that under high water levels and long wave periods, within the flexible *Puccinellia* canopy the orbital velocity was reduced by 35%, while for the more rigid stems of *Elymus*, no significant changes in orbital velocity were found. Differently, under low water levels, and short wave periods, *Elymus* reduced near bed velocity more than *Puccinellia*. As expected, more flexible stems of *Puccinellia* were able to more easily survive the more severe conditions, while the more stiff *Elymus* plants were subject to structural damage.

#### **4. Storms impact on salt marsh morphology**

In comparison to other wetlands, and from a morphological point of view, salt marshes have been found to be more resistant to the impact of storms; this has been mainly attributed to the increased shear strength conferred to the soil by the presence of root systems which are deeper than in other coastal areas such as freshwater wetlands, and floating marshes (e.g. Morton and Barras, 2011). Nevertheless, the impact of storms on salt marshes can significantly vary depending on both storms and ecosystem properties, and can translate into various geomorphic signatures. Some of these signatures have contrasting effects in relation to the long term resilience of the ecosystem. Apart from erosion and deposition processes, affecting marsh platform, marsh shoreline, as well as surrounding tidal flats, storms can also deform the marsh surface through subsurface processes, and incision (e.g. Morton and Barras, 2011). This section presents a summary of some of the main geomorphic impacts of storms on salt marsh ecosystems (Figure 5).

##### **4.1 Incision**

For salt marshes, ponds generated during storms are generally much smaller and less frequent with respect to brackish and freshwater marsh ponds; they also maintain a more amorphous shape (with no preferential direction) in comparison to the more elongated ponds frequently found in freshwater marshes (e.g. Barras, 2011). These ponds are more easily

formed where the terrain is already lower, and strong wind driven currents can erode surface sediments (e.g. Morton et al., 2011). Ponds can then enlarge in time due to subsequent storms, and can also deepen leading to a loss of sediments from the marsh (e.g. Mariotti, 2016). In fact, once the ponds are formed, these can expand even if the rest of the marsh platform is able to keep pace with sea level, and wave action; enlarged ponds can eventually connect to tidal channels (e.g. Mariotti and Fagherazzi, 2013; Schepers et al., 2017).

When a pond is connected to channels, it can recover if its bed is higher than the limit for vegetation growth, or if the deposition rate is larger than the rate of sea level rise. When these conditions are not satisfied, the pond enlarges, becomes susceptible to edge erosion due to internally generated wind waves, and the eroded sediments can get lost through tidal channels (Mariotti and Fagherazzi, 2013). Therefore, depending on the action of biological processes, and sedimentation rates, the formation and enlargement of ponds can be irreversible, or reversible with ponds eventually recovering back to the surrounding marsh platform elevation (e.g. Mariotti and Carr, 2014; Mariotti, 2016).

Plucked marsh features (e.g. Barras et al., 2007) are erosional signatures consisting of irregular scours ranging from around 2 to 20 m which can be found in saline as well as intermediate or freshwater marshes when the mineral matter represents a high percentage of the substrate. Plucked marsh features can occur independently from the elevation with respect to mean sea level, as long as the shear stress is sufficient to incise the areas (e.g. Barras et al., 2007).

## **4.2 Erosion – surface erosion, and lateral erosion**

The denudation of the marsh from the vegetation cover (also referred to as root scalping, e.g. Priestas et al., 2015) can affect areas of the order of kilometres, and occurs when currents and waves induced shear stress strip vegetated surfaces. The depth of denudation determines

the chances and the rate of recovery of the affected areas. If the eroded areas remain above the permanent submerged location, and the root system is not completely destroyed, the denuded zones can recover during the following growing seasons, otherwise the denuded areas might convert to pond or bare tidal flats (e.g. Hendrickson, 1997). The erosion depth of the marsh platform can range from a few to several centimetres. For instance, Hendrickson, 1997, reported erosion rates of 6 cm after the occurrence of Hurricane Erin, and Opal, 1995 for salt marshes in St. Marks River, Florida. However, the erosion of the marsh surface doesn't necessarily correspond to an elevation change as the deformation of the marsh platform through subsurface processes, like compaction or soil swelling, can play an important role as well.

As a consequence of waves generated shear stress, the tidal flats in front of the marsh can deepen which indirectly impacts salt marsh survival, because of an increased depth in front of the marsh can increase wave energy and promote lateral erosion (e.g. Fagherazzi et al., 2006).

The lateral erosion of marsh shorelines has been found to be mainly dictated by the action of wind waves (e.g. Schwimmer, 2001; Marani et al., 2011; Leonardi et al., 2016a, b). For freshwater marshes, the lateral erosion during hurricanes can be up to 100s m; for salt marshes, while wave-induced lateral erosion is in the long term one of the main causes of deterioration, the lateral retreat occurring during hurricanes is relatively low due to the short, and impulsive nature of these events (e.g. Leonardi et al 2016a, b; Figure 6a). Based on a global dataset of salt marsh lateral erosion, and wave data it was found that the yearly retreat rate of marsh shorelines linearly increases with wave energy and a critical threshold in wave energy above which salt marsh erosion drastically accelerates is absent. Such critical threshold is instead more commonly found in sandy environments where erosion drastically increases once the sand dunes are over-washed. While the impact of hurricanes on salt marshes can be very strong, their low frequency and short duration lead to a relatively small effect and they

contribute to only 1% of the erosion in the long term. On the contrary, moderate and frequently occurring storms with a monthly reoccurrence are the most dangerous for salt marsh survival (Leonardi et al., 2016a). It is then reasonable to assume that a storm impacting a stretch of shoreline at 90 degrees has a potential to erode salt marshes which is higher than a storm whose waves are parallel to the shore (e.g. Tonelli et al., 2010).

Finally, in regard to lateral shorelines dynamics, the intensity of wind waves has been found to also modify the shape of marsh boundaries; Leonardi and Fagherazzi, 2014, 2015 showed that the interplay between waves intensity and the spatial variability in marsh resistance determines the shape of marsh shorelines, as well as erosion rates predictability. The variability in erosional resistance is due to the presence of natural heterogeneities caused by different soil resistance and by the variety of ecological, and biological processes interesting different marsh portions. In case of low wave energy conditions, the presence of a variability in erosional resistance might lead to the unpredictable failure of large marsh portions with respect to average erosion rates, and to rough, and jagged marsh boundary profiles displaying high sinuosity values (e.g. Figure 6b, top panel). High-wave-energy conditions, while overall leading to a faster marsh deterioration, cause a constant and predictable erosion, and a smooth marsh boundary profile. A high occurrence of extreme events significantly smooths the marsh boundary, even if it doesn't strongly alter average erosion rates (Figure 6b). Finally, salt marshes subject to weak wave energy conditions are the most susceptible to variations in the frequency of extreme events (Leonardi et al., 2014, 2015).

### **4.3 Deposition**

The occurrence of storms and hurricanes can be accompanied by the deposition of large amount of sediments. As an example, Hurricane Rita generated 4-5 m of storm surge, which resulted in a deposit 0.5m thick, and extending 500 m inland (e.g. Williams, 2009). Cahoon,

2003, 2006 presented a comprehensive set of measurements in regard to elevation changes following the impact of hurricanes at ten sites in the United States; he found deposition rates ranging from a few cm (e.g. 3 cm after Hurricane Emily, 1993, and Gordon, 1994 for salt marshes in North Carolina), up to around 30 cm (e.g. 28, and 20 cm after Hurricane Andrew, 1992, for salt marshes in Bayou Chitigue, and Old Oyster, Louisiana).

Depending on the net direction of sediment transport, deposits may be laid down over the salt marsh surface or translated seaward. Storms may not, therefore, necessarily leave behind distinct depositional units but instead increase the increment of tidal deposition through elevated suspended sediment concentrations and/or flow velocities (Stumpf, 1983), thus enhancing the usual mechanisms of settling during inundations or over-bank spilling in close proximity to creeks or the point of tidal ingress. Indeed, Turner et al. (2006) suggest that large storms increase the supply of mineral matter from offshore via tidal creeks, and have shown that, for Mississippi River salt marshes, the density of minerogenic sediments in salt marsh cores increases in concert with the occurrence of major hurricanes (Turner et al., 2007).

Deposition during storms is readily evidenced where breaching and flooding of the supratidal coastline occurs, e.g. washover deposits or fans. For example, Scileppi and Donnelly (2007) found that washover deposits on the Long Island coast correlate with landfalls of the most intense documented hurricanes, and that periods of increased and decreased landfall incidence can be evidenced in the back-barrier sediment record (cf. Liu and Fearn, 2000; Donnelly et al., 2001; 2004). Barrier overwashing during storms can also deposit lobes of sand and intermixed shells over back-barrier salt marshes, where shell beds may then be preserved in the sediment record as an archive of storm washover (Ehlers et al., 1993). Extensive washover deposits resulting from storms have also been found in a back-barrier setting along the Chenier Plain of Louisiana where the intensity of recent hurricanes influences the extent and grain size of the deposits (Williams, 2011).

It is less common for salt marshes to preserve depositional evidence of storms, or at least deposits that can readily be distinguished from the usual background of regular tidal deposition or, indeed, other extreme events such as tsunami (cf. Goff et al., 2004; Morton et al., 2007). Goodbred and Hine (1993) recorded the deposition of a tan to grey unit of clays, silt to very fine sand, and marine biogenic matter across Waccasassa Bay salt marshes in Florida following a 3 m storm surge. The deposit was made up of sedimentary material similar to that of the underlying marsh sediments, indicating a local origin. Proximity to tidal ingress had a significant influence on the thickness of the deposit, increasing from a few cm on the salt marsh surface to as much as 12 cm along creek margins. Generally, severe storms have the potential to deposit distinctive sand units that thin and fine in a landward direction over 100s of meters, that have a sharp basal contact with the underlying salt marsh deposits, and that contain marine microfossils (e.g. Morton and Sallenger, 2003; Turner et al., 2006; Williams, 2009). Such anomalous deposits are characterized using several criteria such as the extent of inundation, landward-thinning and/or landward-fining of the deposit, single or multiple particle size grading, and contained microfossil assemblage (Hawkes and Horton, 2012).

Similar, unconformable sand deposits can be found within the salt marsh sediment record of back-barrier estuaries along the Central Coast of California (e.g. Clarke et al., 2014) where their incidence is connected to barrier breaching and inundation during storms. In this case, high frequency variability in the particle size of such deposits in the back-barrier stratigraphy can be associated with ENSO-driven storms, but where the barrier breaching is most likely due to high river flow as opposed to coastal erosion during storms (Clarke et al., 2017).

Drawing on examples from the longer Holocene sediment record, Haggart (1988) examined the stratigraphic and sedimentary evidence of a tidal surge deposit in two open estuary settings in north-eastern Scotland. This micaceous, silty sand was deposited across

pre-existing inter-tidal to perimarine environments, which then returned immediately following its deposition. The stratigraphic evidence is therefore indicative of a high energy environment affecting a wide range of coastal environments simultaneously, with a vertical range of 3.5-5.0 m. Detailed dating, particle size, and paleoecological data reveal this deposit to be marine in origin and virtually instantaneous in its deposition. Similar deposits of this kind are found in a number of estuarine and back-barrier settings in north-east Scotland (Smith et al., 2004) for which the timing, rarity, and run-up (as much as 25 m) are indicative of a tsunami rather than a storm surge. Information on storm-related sediment redistribution across the salt marsh surface can equally come from evidence other than stratigraphic, grain size or palaeoecological data. For example, Rahman et al. (2013) explored down-core trends in radioactive pollution to determine patterns of sedimentation in north-west England. A secondary increase in both  $^{241}\text{Am}$  and  $^{137}\text{Cs}$  activity in the upper 5-10 cm of salt marsh cores from the Dee was interpreted as the re-deposition of sediments eroded from the salt marsh edge, linked to a severe storm in 1990. In principle, the erosion and redistribution of historical pollutants in industrialized estuaries can also be revealed by the analysis of heavy metals or persistent organic pollutants.

In summary, storm deposits are more readily apparent in back-barrier salt marshes where coastal breaching and overwashing enable the landward penetration of coarse sediment lobes that then appear anomalous against the background of tidal mud deposition. Such deposits also have the potential to be found in more open estuary settings where the storm surge results in the landward transport of coarse marine sediment or increases the potential for the redistribution of eroded material onto the salt marsh surface. Identifying such deposits requires a multi-proxy approach to evidence not only the nature and dynamics of the depositional environment but also the age and origin of the sediments, particularly for reconstructing periods of increased and decreased storminess.

#### **4.4 Deformation**

Apart from surface processes of erosion, deposition, and incision, subsurface processes induced by soil compaction or groundwater flow are also an important consequence of storms and storm surges occurrence, and can lead to substantial deformation or changes in marsh elevation.

Soil compaction due to storm surge water is quite common; for instance, after hurricane Andrew, 1992, and for salt marshes in Bayou Chitigue, Louisiana, in spite of a 28 cm thick deposit, the total change in elevation was -5cm due to soil compaction (Cahoon, 2006). Similarly, for salt marshes in Cedar Island, North Carolina, the surface erosion due to Hurricane Felix, and Jerry was only -1cm, but the change in elevation due to soil compaction reached -18cm (Cahoon et al., 1999; Cahoon, 2006). Soil shrinkage or swell can be also caused by an alteration of water fluxes mainly induced by storm surge events. According to Hendrickson, 1997, soil shrinkage caused a 13 cm, and 8 cm lowering of the marsh platform for salt marshes in Florida after Hurricane Opal, 1995 and Erin, 1995 respectively. On the contrary, during Hurricane Alberto, 1994, soil swelling caused by the storm surge increase in water content, caused an increase in elevation of 13 cm for the salt marshes in Florida, (Cahoon, 2006).

## **5. Storms impact on salt marsh sediment budget**

A salt marsh is defined not only through the vegetated marsh plain, but by the entire geomorphic complex. This complex includes the adjacent estuarine/marine seabed, tidal marsh channels, intertidal flats, marsh scarps, the marsh plain, and pools within the marsh plain. Though the salt marsh plain can accrete vertically through organic and inorganic sediment accretion, the geomorphic evolution of the other components is influenced by the inorganic sediment budget (e.g. Ganju et al., 2017).



Sources of sediment for coastal salt marshes are diverse, but can broadly be categorized into external sources, from the erosion of neighbouring coasts or seafloor and from riverine sediment discharge, as well as internal sources from sediment resuspension on intertidal mudflats adjacent to the salt marshes or erosion of the marsh edges and tidal channels (Schuerch et al., 2014). All sources can be highly variable in time and space and are often driven by highly energetic events, such as storms causing severe precipitation, storm surges and/or wave setup (Ma et al., 2014; Schuerch et al., 2016).

The transport of sediments to the salt marsh occurs on multiple timescales. Wind-waves, due to diurnal or stronger episodic winds, can mobilize estuarine and intertidal flat sediments, erode marsh scarps, and increase sediment concentrations in the water column (Fagherazzi and Priestas, 2010; Ganju et al. 2013).

Over large and small spatio-temporal scales, the net sediment budget will govern whether the complex is trending towards expansion or contraction. For example, a sediment transport deficit that results in a deepening of the estuary will allow for greater propagation of wave energy towards the marsh scarp, leading to increased thrust and erosion of the scarp. The sediment liberated from the marsh scarp may then deposit elsewhere in the complex, or it may be exported from the entire system through hydrodynamic processes. Inorganic sediment supply is also important for vertical accretion on marsh plains (Reed 1989), though in some environments marshes can subsist entirely on organic production (Turner et al. 2002). Furthermore, where the marsh plain meets the marsh scarp, there is a more delicate balance that is dependent on sediment supply, and morphological features as well; for instance, Redfield (1972) identifies the tendency for slumped blocks of peat to trap sediment, and reconstitute marsh plain through recolonization by vegetation, thereby leading to no net loss of marsh plain.

Storms can have varying effects on sediment supply: in some cases they lead to massive sediment export from the system (Ganju et al. 2013), substantial sediment import (Rosencranz et al. 2016), significant marsh plain deposition (Goodbred and Hine, 1995), or negligible marsh plain deposition (Elsey-Quirk 2106).

Ganju et al. (2013) identified disparate sediment sources and transport mechanisms at two Chesapeake Bay marsh complexes (one stable, one degraded), i.e., tidal processes delivered sediment to the stable marsh while fall and winter storms exported sediment from the degraded marsh. Conversely, Rosencranz et al. (2016) found that a single 3 day storm delivered enough sediment to counteract two months of tidally driven sediment export within a Pacific coast marsh complex.

For a degraded marsh complex in Blackwater, MD, USA, tidal resuspension and advection did not provide sediments, while sustained northwest wind events with a 2-wk return interval were able to both mobilize sediment from open-water areas and export sediments (Ganju et al., 2013, Figure 7b); the orientation of the open-water area was aligned along the northwest-southeast axis, thereby allowing for greater fetch and wind-wave exposure during northwest winds. The ensuing wind-waves both mobilized subaqueous sediments and eroded marsh edges; export was then caused by a regional hydrodynamic response which led to net water export. However, a nearby stable complex (Fishing Bay, MD, USA, Figure 7a) imported sediment due to tidal resuspension/advection and proximity to an estuarine sediment source. There was minimal sediment export during the same aforementioned wind-wave events, due to a lack of open-water area.

In Barnegat Bay, New Jersey (USA) a strong south-to-north gradient in shoreline type and sediment availability leads to a variable response to storm events. Dinner Creek, in the southern portion of the bay, is bordered by undeveloped marsh shoreline and shoals consisting of fine sediment (Miselis et al. 2016; Ganju et al. 2014), while Reedy Creek is surrounded by

615 hardened shorelines and coarse-sediment dominated shoals. Ganju et al. (2017) reported a net  
616 sediment import for Dinner Creek and negligible sediment transport in Reedy Creek;  
617 cumulative fluxes in response to wind events indicate a direction-dependent response (Figure  
618 7c, d). Both sites export sediment during periods with northwest winds and import sediment  
619 during southerly winds, but Dinner Creek imports sediment during easterly winds while Reedy  
620 Creek remains neutral (Figure 7c, d). This differential response is likely due to the availability  
621 of sediment in the estuary. These results show that the location of a salt marsh plays a strong  
622 role in the sediment dynamics during storm events, with varied directional responses. Tidal  
623 asymmetry affects the net import/ export of sediments as well. The distortion of the tidal wave  
624 may significantly change under storm conditions, hence converting a system which would  
625 normally import sediments into a system which export sediments (Schuerch et al., 2014).

626 Finally, Ganju et al. (2017) synthesized sediment budgets of eight microtidal salt marsh  
627 complexes, and demonstrated a relationship between the sediment budget and the unvegetated-  
628 vegetated marsh ratio, indicating that sediment deficits are linked to conversion of vegetated  
629 marsh portions to open water. Both observational and modelling efforts provide insight into  
630 the influence of storms and extreme events on sediment transport to and from salt marshes.

## 631 632 **Storms impact on sea level rise resilience**

633 Accelerated sea level rise is challenging the survival of coastal salt marshes, which may  
634 decrease in elevation within the tidal frame and eventually be inundated too frequently to  
635 support the growth of salt marsh vegetation (Kearney et al., 1988; Day et al., 2000; Schepers  
636 et al., 2017). With increasing rates of sea level rise, coastal salt marshes rely on a higher  
637 sediment supply in order to vertically adapt to the rising sea level (French, 1993; Kirwan et al.,  
638 2010a; D'Alpaos et al., 2011). Ma et al. (2014), for example, show a decrease in marsh  
639 sedimentation rates in the Oosterschelde estuary (NL) after the construction of a storm surge

barrier, which markedly reduced the (external) marine sediment delivery, but also show that sedimentation rates are still keeping up with sea level rise due to sediment resuspension on the adjacent intertidal mudflat during storm events.

Although estimates of critical rates of sea level rise for coastal salt marshes around the world indicate a relatively high resilience for many salt marsh sites (Kirwan et al., 2016), all assessments also highlight that the available sediment supply is a key factor for marsh resilience to sea level rise (French, 2006; Kirwan et al., 2010a; D'Alpaos et al., 2011; Schuerch et al., 2013). Furthermore, salt marshes in microtidal regimes were identified as particularly sensitive to a drop in sediment supply under increasing rates of sea level rise, whereas salt marshes in macrotidal regimes are more resilient to high rates of sea level rise and/or reduced sediment supply (Spencer et al., 2016; Kirwan et al., 2010b). While being more susceptible to drowning as a consequence of sea level rise, sedimentation rates on microtidal marshes were also shown to be more responsive to changes in storm activity due to an increase in sediment supply through intertidal sediment resuspension with respect to macrotidal marshes. Kolker et al. (2009), for example, found clear storm signals in the sedimentation records of their microtidal and wave exposed study sites within the Long Island Sound (USA), but a much reduced signal in the neighbouring macrotidal sites.

In this context, elongated periods (decades) of increased storm activity appear as the main driver for sedimentation in excess of local sea level rise rates as shown for a mesotidal salt marsh in the German North Sea (Figure 8; Schuerch et al., 2012). This excess sedimentation significantly contributes to the resilience of the marsh with respect to its vertical performance and its ability to adapt the future SLR (Schuerch et al., 2013). In the Mississippi Delta, extreme events such as the hurricanes Katrina and Rita in 2005 were reported to contribute sediment layers of 9-13 and 7 cm, respectively, which is a manifold of the regular annual sedimentation (Horton et al., 2009). Meanwhile, Tweel and Turner (2014) argue that

the strongest 2% of extreme events contribute 15% of the sedimentation to the marshes of the Mississippi Delta, whereas the majority of the sedimentation (78%) can be attributed to moderate hurricanes with a landfall barometric pressure between 930 and 960 mb (Tweel and Turner, 2014). In addition to sediment deposition, subsurface processes may, however, dominate the elevation response to storm events in many marshes of the Mississippi Delta (Cahoon, 2006; McKee and Cherry, 2009). Subsurface processes are primarily related to soil organic matter, hence are most relevant in organogenic marshes and less so in minerogenic marshes.

Moderate storm events also appear to be responsible for the majority of marsh sedimentation on the Danish peninsula of Skallingen (Bartholdy et al., 2004), where extreme storm events were shown to increase suspended sediment concentrations within the adjacent tidal basin by a factor of up to 20 due to sediment resuspension on the intertidal mudflats. There, a single extreme event could contribute 7.5% to the annual sediment deposition, whereas a single regularly occurring gale already contributes 71% of this (Bartholdy and Aagard, 2001). The high importance of frequently inundating gale events is in accordance with the modelling study of Schuerch et al. (2013), who suggest that the frequency of storm events is more important for inorganic marsh accretion than their intensity. The explanation for this behaviour is that the frequency distribution of high and extreme water levels decreases exponentially with increasing high water levels (Bartholdy et al., 2004; Schuerch et al., 2013), whereas the sediment resuspension on the intertidal mudflat appears to follow a linear relationship with increasing high water level (Temmerman et al., 2003) or significant wave heights (Fagherazzi and Pristas, 2010). Therefore extreme sediment resuspension events are too rare to make a significant impact. Furthermore, the impact of wave-induced sediment resuspension decreases with increasing water depths during high inundation events (Fagherazzi and Wiberg, 2009; Christiansen et al., 2006).

However, sediment resuspension within the intertidal zone is a highly variable process (Carniello et al., 2016), as it also relies on the sediment composition of the seabed and the presence of benthic biology determining the erosion thresholds and the stability of the seabed (Le Hir et al., 2007; Grabowski et al., 2011). In particular the benthic biological activity (e.g. vegetated seabeds, diatom biofilms, and benthic macrofauna) has the potential to introduce significant spatial and temporal variations in sediment resuspension (Andersen et al., 2001). Locally, and depending on biological activity, the impact of storm events on the sediment supply of coastal salt marshes may therefore be subject to considerable seasonal variations, often with a stronger impact of storm events on sediment supply during the winter months (Temmerman et al., 2003).

During elongated periods of increased storm activity, which appear to be most effective in increasing sedimentation rates on salt marshes (Figure 8; Schuerch et al., 2012), intertidal sediment resuspension may cause a lowering of the mudflat elevation and potentially conversion to a subtidal flat. In combination with an enhanced vertical growth of the vegetated marsh platform this may lead to an increased mudflat-salt marsh elevation gradient (Le Hir et al., 2007; Mariotti and Fagherazzi, 2010). Incoming waves, therefore, have an increased erosive impact on the steeper marsh edge, hence increasing the marsh's vulnerability to lateral erosion (e.g. Van de Koppel et al., 2005)). A reduction of the intertidal mudflat area due to storm erosion also reduces the sediment resuspension and therefore the sediment supply for the vertical growth of the salt marsh. Both marsh edge erosion and the vertical performance of coastal salt marshes are therefore critically dependent on external sediment supply, which in fact is often enhanced by storm events as well (Mariotti and Carr, 2014).

The sediment import into the tidal basins of the Wadden Sea (South-eastern North Sea), for example, increases during storm events and the sediment composition shifts into the coarser spectrum as increased erosion takes place along the beaches of the adjacent barrier islands and

the ebb-tidal delta (Schuerch et al., 2014). Similarly, increased suspended sediment concentrations are observed along the UK East coast as a consequence of the erosion of soft cliffs, particularly during the winter season and intensified storm periods (McCave, 1987; Nicholls et al., 2000; Dyer and Moffat, 1998). Storm events are also often associated with increased precipitation in the catchments of the rivers draining into the coastal zone. The increased river runoff often increases the sediment delivery into the coastal zone and hence the “external” sediment supply for coastal salt marshes (Schuerch et al., 2016). The relationship between river runoff and sediment delivery is, however, not necessarily a straightforward one as it is subject to intense anthropogenic modifications, such as river damming or land use change in the river catchment (Syvitski et al., 2005).

Despite the abundant field evidence and the well-developed knowledge on the importance of sediment supply for coastal salt marshes, current estimations of future salt marsh development largely neglects the processes and feedbacks involved in storm-related sedimentation by neglecting the temporal variations in sediment supply and assuming a constant sediment supply throughout the coming century (e.g. Kirwan et al., 2010; D’Alpaos et al., 2011; Mariotti and Carr, 2014). Accounting for the storm-induced variability in sediment supply for coastal salt marshes in future models is particularly important as storm activity is known to be subject to significant decadal variability (e.g. driven by the North-Atlantic Oscillation) and may prevent or facilitate the collapse of coastal salt marshes, when conventional modelling under the assumption of constant sediment supply and storm activity would predict differently.

## **Discussion and Conclusions**

In face of climate change, the continued delivery of salt marsh ecosystem services, such as mitigation of flood risks and shoreline erosion risks, and carbon sequestration, is increasingly important.

Under storm conditions salt marshes are able to effectively dissipate both high water levels and wave energy even under extreme water level conditions such as during storm surges, and even if the wave-bottom interaction, and energy dissipation decreases with increasing water level. Empirical data and modelling studies demonstrate effective storm surge height reduction behind large (at least 10 km wide) and continuous marshes during moderate storm surges, but also point at limitations in the storm surge protection value, when marshes are smaller, intersected by large channels or open water areas, and during extreme storm surges. This implies that storm surge protection schemes should ideally rely on a combination of conservation and restoration of large continuous marsh areas, where space is available, and engineered flood defences, where necessary (Temmerman et al. 2013).

Under storm surge conditions, up to 60% of the wave attenuation is attributable to the sole presence of vegetation, rather than to the decrease in water depth on the marsh platform relative to the surrounding tidal flat (e.g. Moeller et al., 2014). Vegetation properties largely influence this dissipation process; while the more flexible stems tend to flatten during powerful storms (with a reduction in dissipation potential), they are also the more resilient to structural damage, and their flattening helps to protect the marsh substrate against erosion. On the other hand, with increasing wave energy, high vegetation stiffness can enhance the turbulence and surface erosion around plant stems (Silinski et al., 2016; Rupprecht et al., 2017).

Storm action can have various impacts on the geomorphological evolution of salt marshes, and different implications for their long term survival to sea level rise, and climate change in general. Storms impact potentially causes erosion of marsh boundaries, marsh platforms, and surrounding tidal flats, but it might also deliver substantial amount of sediments to the marsh platform.



Under the assumption of an increase in magnitude, and reduced frequency of extreme events it can be argued that the after-storm impact on marsh boundaries is expected to be only slightly affected by such changes; this is because it has been shown that the lateral erosion of salt marshes is mostly dictated by average weather conditions rather than by extreme events. The biggest impact that storms could have in relation to lateral salt marsh dynamics could instead be connected to the deepening of tidal flats which promote higher wave energy at the marsh boundary, and reduces wave energy dissipation by bottom friction, causing therefore an increase in the erosion potential during inter-storms period, i.e. under normal weather conditions.

The impact on the vertical salt marsh dynamic is complicated because, even if more intense storms have the potential to deposit more sediments, there are evidences about the fact that storms frequency is more important than intensity for the long term inorganic accretion of salt marshes. The explanation for this behaviour is that the frequency distribution of high and extreme water levels decreases exponentially with increasing high water levels (Schuerch et al., 2013, 2014).

The occurrence of storms might then directly or indirectly impact the sediment budget of the coastline. In particular, the direction of storm events can determine whether there is a direct import or export from a coastal embayment. Furthermore, the occurrence of storms is generally connected to precipitation events and surface runoff which might increase the transport of sediments from the catchment to the coastline (e.g. Ganju et al., 2013)

The latter considerations highlight the necessity to focus on the indirect impact that large storms might exert on salt marshes not only in the immediate after storm period, but also in the longer term, and on how their morphological consequences influence the response of the system to normal weather conditions during inter-storm periods. Some of the challenges

highlighted from the complexity of the problem also include the necessity to consider salt marsh systems as a whole by adopting an integrated approach, taking into account the marsh tidal flat continuum and by accounting for various sediment sources.

## Figures

### Figure 1

Percentage changes in Emmanuel's (1995) wind maximum potential intensity ( $MPI_v$ ) per degree increase in global surface air temperature. Large values of  $MPI_v$  values are generally associate to enhanced tropical storms activity, and intensity (adapted from Vecchi and Soden, 2007).

### Figure 2

Sketch of mechanisms and sediment fluxes possibly responsible for salt marsh vertical and horizontal dynamics. Black dashed box represents an hypothetical control volume for the evaluation of the sediment budget.

### Figure 3

Relationship between the attenuation rate of High Water Levels ( $dH_{WL}/dx$ ) at least 0.4m above the marsh platform, and  $\alpha_v$ , i.e. ratio between the over-marsh water volume ( $V_{pl}$ ) and

the total water volume ( $V_{pl}+V_c$ , i.e. over-marsh water volume + water volume within channels) (adapted from Stark et al., 2016).

#### **Figure 4**

Reduction of total Energy [ $J\ m^{-2}$ ] between sand flat, marsh edge and marsh interior for ten representative measurements ‘bursts’ (adapted from Moeller, 1999).

#### **Figure 5**

Diagram representative for some of the major morphologic impacts of storms on salt marshes, their spatial scale, and useful literature references. Morton and Barras, 2011; b) Mariotti and Carr, 2014; c) Mariotti, 2016; d) Fan et al., 2006; e) Scileppi and Donnelly, 2007; f) Williams, 2009; g) Leonardi et al., 2016a,b; h) Leonardi et al., 2014, 2015; i) Barras, 2007, l) Cahoon, 2006; m) Cahoon, 2003; These impact are mainly categorized into the following: Deformation, Erosion, Deposition, and Incision.

#### **Figure 6**

A) Contribution of different wind categories to salt marsh erosion (from Leonardi et al., 2016). B) Impact of increasing extreme events frequency on the shape of marsh shorelines (adapted from Leonardi et al., 2014, 2015). Increasing the occurrence of extreme events smooths the marsh shoreline.

#### **Figure 7**

Sediment flux response to wind forcing at four wetland complexes, as a function of wind direction (radial position) and speed (outward position). The wind direction indicates direction the wind is coming from. Fishing Bay and Blackwater (Maryland, USA), are adjacent to Chesapeake Bay, but their respective locations relative to sediment sources and external forcing result in disparate sediment transport responses to wind events. Northwest winds export sediment from both sites, but southerly winds allow for sediment import at Fishing Bay due to proximity to a southern sediment source (Ganju et al., 2013). Dinner and Reedy Creeks, in southern and northern Barnegat Bay (New Jersey, USA), respectively, both export sediment during westerly winds, but Dinner Creek imports sediment during strong easterly winds. This is likely due to increased fine sediment availability and undeveloped shoreline in the southern portion of Barnegat Bay, as opposed to coarser sediments and hardened shoreline in northern Barnegat Bay.

## **Figure 8**

(a) Historic marsh elevations in comparison to the development of the mean high water level (MHW) and the mean sea level (MSL) for three cores (S1: high marsh; S2: low marsh; S3: pioneer marsh) from a salt marsh on the German island of Sylt (in the South-eastern North Sea). Deposition dates were derived from  $^{210}\text{Pb}$  and  $^{137}\text{Cs}$  data (open diamonds). The green shaded area indicates the periods of excess sedimentation during periods of increased storm activity. (b) Comparison of sedimentation rates (stars) at core location S2 with storm frequency (open circles), defined as the number of water levels exceeding 2.4 m above the long-term mean sea level (NN: German ordnance datum). Modified after Schuerch et al. (2012).

## **Acknowledgments**

854 The following funding are acknowledged: USGS Federal award No G16AC00455; NERC  
855 NE/N015614/1.

## 856 **References**

857 Andersen, T.J., 2001. Seasonal Variation in Erodibility of Two Temperate, Microtidal  
858 Mudflats. *Estuarine, Coastal and Shelf Science*, 53(1), 1-12.

859 Augustin, L.N., Irish, J.L., Lynett, P., 2009. Laboratory and numerical studies of wave damping  
860 by emergent and near-emergent wetland vegetation. *Coastal Engineering* 56, 332–  
861 340.

862 Bacmeister, J.T., Reed, K.A., Hannay, C., Lawrence, P., Bates, S., Truesdale, J.E.,  
863 Rosenbloom, N. and Levy, M., 2016. Projected changes in tropical cyclone activity  
864 under future warming scenarios using a high-resolution climate model. *Climatic*  
865 *Change*, pp.1-14.

866 Barbier, E. B., Georgiou, I. Y., Enchelmeyer, B., and Reed, D. J., 2013, The Value of Wetlands  
867 in Protecting Southeast Louisiana from Hurricane Storm Surges Plos One, v. 8, p.  
868 e58715.

869 Barras, J.A., 2007. *Satellite images and aerial photographs of the effects of Hurricanes Katrina*  
870 *and Rita on coastal Louisiana* (No. 281). Geological Survey (US).

871 Bartholdy, J. and Aagaard, T., 2001. Storm surge effects on a back-barrier tidal flat of the  
872 Danish Wadden Sea. *Geo-Marine Letters*, 20(3), 133-141.

873 Bartholdy, J., Christiansen, C. and Kunzendorf, H., 2004. Long term variations in backbarrier  
874 salt marsh deposition on the Skallingen peninsula - the Danish Wadden Sea. *Marine*  
875 *Geology*, 203(1-2), 1-21.

876 Bouma, T. J., De Vries, M. B., Low, E., Peralta, G., Tanczos, I., Van de Koppel, J., and  
877 Herman, P. M. J., 2005, Trade-offs related to ecosystem-engineering: a case study on  
878 stiffness of emerging macrophytes: *Ecology*, v. 86, p. 2187-2199.

879 Cahoon, D.R., Day Jr, J.W. and Reed, D.J., 1999. The influence of surface and shallow  
880 subsurface soil processes on wetland elevation: A synthesis. *Current topics in wetland*  
881 *biogeochemistry*, 3, pp.72-88.

882 Cahoon, D. R. 2003. Storms as agents of wetland elevation change: their impact on surface and  
883 subsurface sediment processes, Proceedings of the International Conference on Coastal  
884 Sediments 2003. May 18-23, 2003, Clearwater Beach, FL, USA. CD-ROM Published  
885 by World Scientific Publishing Corp. and East Meets West Productions, Corpus Christi,  
886 Texas, USA. ISBN 981-238-422-7.

887 Cahoon, D.R., 2006. A review of major storm impacts on coastal wetland elevations. *Estuaries*  
888 *and Coasts*, 29(6), pp.889-898.

889 Carniello, L., D'Alpaos, A., Botter, G. and Rinaldo, A.C.J.F., 2016. Statistical characterization  
890 of spatiotemporal sediment dynamics in the Venice lagoon. *Journal of Geophysical*  
891 *Research: Earth Surface*, 121(5), 1049-1064.

892 Cheong, S. M., Silliman, B., Wong, P. P., van Wesenbeeck, B., Kim, C. K., and Guannel, G.,  
893 2013, Coastal adaptation with ecological engineering: *Nature Climate Change*, v. 3, no.  
894 9, p. 787-791.

895 Christiansen, T., Wiberg, P.L. and Milligan, T.G., 2000. Flow and sediment transport on a tidal  
896 salt marsh surface. *Estuarine, Coastal and Shelf Science*, 50(3), pp.315-331.

---

897 Christiansen, C., Vølund, G., Lund-Hansen, L.C. and Bartholdy, J., 2006. Wind influence on  
898 tidal flat sediment dynamics: Field investigations in the Ho Bugt, Danish Wadden Sea.  
899 *Marine Geology*, 235(1-4), 75-86.

900 Clarke, D. W., Boyle, J. F., Plater, A. J. (2017). Particle-size evidence of barrier estuary regime  
901 as a new proxy for ENSO climate variability. *Earth Surface Processes and Landforms*,  
902 *in press*. doi:10.1002/esp.4106

903 Clarke, D. W., Boyle, J. F., Chiverrell, R. C., Lario, J., Plater, A. J. (2014). A sediment record  
 904 of barrier estuary behaviour at the mesoscale: Interpreting high-resolution particle size  
 905 analysis. *Geomorphology* 221, 51-68.

906 Costanza, R., Perez-Maqueo, O., Martinez, M. L., Sutton, P., Anderson, S. J., and Mulder, K.,  
 907 2008, The value of coastal wetlands for hurricane protection: *Ambio*, v. 37, no. 4, p.  
 908 241-248.

909 CPRA, 2012, Louisiana's Comprehensive Master Plan for a Sustainable Coast: Louisiana  
 910 Coastal Protection and Restoration Authority.

911 D'Alpaos, A., Mudd, S.M. and Carniello, L., 2011. Dynamic response of marshes to  
 912 perturbations in suspended sediment concentrations and rates of relative sea level rise.  
 913 *Journal of Geophysical Research-Earth Surface*, 116(F4), F04020.

914 D'Alpaos, A. and Marani, M., 2016. Reading the signatures of biologic–geomorphic feedbacks  
 915 in salt-marsh landscapes. *Advances in Water Resources*, 93, pp.265-275.

916

917 Danielsen, F., Sorensen, K., Olwig, M. F., Selvam, V., Parish, F., Burgess, N. D., Hiraishi, T.,  
 918 Karunagaran, F. M., Rasmussen, M. S., Hansen, L. B., Quarto, A., and Suryadiputra,  
 919 N., 2005, The Asian Tsunami: a protective role for coastal vegetation: *Science*, v. 310,  
 920 p. 643-643.

921 Day, J., Britsch, L., Hawes, S., Shaffer, G., Reed, D. and Cahoon, D., 2000. Pattern and process  
 922 of land loss in the Mississippi Delta: A Spatial and temporal analysis of wetland habitat  
 923 change. *Estuaries and Coasts*, 23(4), 425-438.

924 Das, S., and Vincent, J. R., 2009, Mangroves protected villages and reduced death toll during  
 925 Indian super cyclone: *Proceedings of the National Academy of Sciences of the United*  
 926 *States of America*, v. 106, p. 7357-7360.

927 Day, J. W., Boesch, D. F., Clairain, E. J., Kemp, G. P., Laska, S. B., Mitsch, W. J., Orth, K.,  
 928 Mashriqui, H., Reed, D. J., Shabman, L., Simenstad, C. A., Streever, B. J., Twilley, R.  
 929 R., Watson, C. C., Wells, J. T., and Whigham, D. F., 2007, Restoration of the  
 930 Mississippi delta: lessons from hurricanes Katrina and Rita: *Science*, v. 315, p. 1679-  
 931 1684.

932 Deegan, L. A., Johnson, D. S., Warren, R. S., Peterson, B. J., Fleeger, J. W., Fagherazzi, S.,  
 933 and Wollheim, W. M., 2012, Coastal eutrophication as a driver of salt marsh loss:  
 934 *Nature*, v. 490, p. 388-392.

935 de Groot, A.V., Veeneklaas, R.M. and Bakker, J.P., 2011. Sand in the salt marsh: Contribution  
 936 of high-energy conditions to salt-marsh accretion. *Marine Geology*, 282(3-4), 240-254.

937 Dyer, K.R. and Moffat, T.J., 1998. Fluxes of suspended matter in the East Anglian plume  
 938 Southern North Sea. *Continental Shelf Research*, 18(11), 1311-1331.

939 Dijkema, K.S., 1987. Geography of salt marshes in Europe. *Zeitschrift für Geomorphologie*  
 940 N.F. 31 (4), 489-499.

941 Donnelly, J. P., Roll, S., Wengren, M., Butler, J., Webb III, T. (2001) Sedimentary evidence  
 942 of intense hurricane strikes from New Jersey. *Geology* 29, 615–618.

943 Donnelly, J. P., Butler, J., Roll, S., Wengren, M., Webb III, T. (2004) A backbarrier overwash  
 944 record of intense storms from Brigantine, New Jersey. *Marine Geology* 210, 107–121.

945 Ehlers, J., Nagorny, K., Schmidt, P., Stieve, B., Zietlow, K. (1993) Storm Surge Deposits in  
 946 North Sea Salt Marshes Dated by  $^{134}\text{Cs}$  and  $^{137}\text{Cs}$  Determination. *Journal of Coastal*  
 947 *Research* 9(3), 698-701.

948 Elsey-Quirk, T., 2016. Impact of Hurricane Sandy on salt marshes of New jersey. *Estuarine,*  
 949 *Coastal and Shelf Science*, 183, pp.235-248.

950 Elwany, M., W. O'Reilly, R. Guza, and R. Flick (1995), Effects of Southern California kelp  
 951 beds on waves, *J. Waterw. Port Coastal Ocean Eng.*, 121(2), 143–150.



952 Fan, D., Guo, Y., Wang, P. and Shi, J.Z., 2006. Cross-shore variations in morphodynamic  
 953 processes of an open-coast mudflat in the Changjiang Delta, China: with an emphasis  
 954 on storm impacts. *Continental Shelf Research*, 26(4), pp.517-538.

955 Fagherazzi, S., Carniello, L., D'Alpaos, L. and Defina, A., 2006. Critical bifurcation of shallow  
 956 microtidal landforms in tidal flats and salt marshes. *Proceedings of the National*  
 957 *Academy of Sciences*, 103(22), pp.8337-8341.

958 Fagherazzi, S. and Wiberg, P.L., 2009. Importance of wind conditions, fetch, and water levels  
 959 on wave-generated shear stresses in shallow intertidal basins. *Journal of Geophysical*  
 960 *Research*, 114(F3), F03022.

961 Fagherazzi, S. and Priestas, A.M., 2010. Sediments and water fluxes in a muddy coastline:  
 962 interplay between waves and tidal channel hydrodynamics. *Earth Surface Processes and*  
 963 *Landforms*, 35(3), 284-293.

964 Fagherazzi, S., 2013. The ephemeral life of a salt marsh. *Geology*, 41(8), pp.943-944.

965 French, J.R., 1993. Numerical simulation of vertical marsh growth and adjustment to  
 966 accelerated sea-level rise, North Norfolk, U.K. *Earth Surface Processes and Landforms*,  
 967 18(1), 63-81.

968 French, J., 2006. Tidal marsh sedimentation and resilience to environmental change:  
 969 Exploratory modelling of tidal, sea-level and sediment supply forcing in predominantly  
 970 allochthonous systems. *Marine Geology*, 235(1-4), 119-136.

971 Ganju, N.K., Defne, Z., Kirwan, M.L., Fagherazzi, S., D'alpaos, A. and Carniello, L., 2017.  
 972 Spatially integrative metrics reveal hidden vulnerability of microtidal salt  
 973 marshes. *Nature communications*, 8.

974 Ganju, N.K., Miselis, J.L. and Aretxabaleta, A.L., 2014. Physical and biogeochemical controls  
 975 on light attenuation in a eutrophic, back-barrier estuary. *Biogeosciences*, 11, pp.7193-  
 976 7205.

977 Ganju, N.K., Nidzieko, N.J. and Kirwan, M.L., 2013. Inferring tidal wetland stability from  
 978 channel sediment fluxes: Observations and a conceptual model. *Journal of Geophysical*  
 979 *Research: Earth Surface*, 118(4), pp.2045-2058.

980 Ganju, N.K., Schoellhamer, D.H. and Bergamaschi, B.A., 2005. Suspended sediment fluxes in  
 981 a tidal wetland: Measurement, controlling factors, and error analysis. *Estuaries and*  
 982 *Coasts*, 28(6), pp.812-822.

983

984 Goff, J., McFadgen, B.G., Chagué-Goff, C. (2004) Sedimentary differences between the 2002  
 985 Easter storm and the 15th-century Okoropunga tsunami, southeastern North Island,  
 986 New Zealand Marine Geology 204(1–2), 235-250.

987 Goodbred, S.L. and Hine, A.C., 1995. Coastal storm deposition: Salt-marsh response to a  
 988 severe extratropical storm, March 1993, west-central Florida. *Geology*, 23(8), pp.679-  
 989 682.

990 Goodbred Jr, S.L., Wright, E.E. and Hine, A.C., 1998. Sea-level change and storm-surge  
 991 deposition in a late Holocene Florida salt marsh. *Journal of Sedimentary Research*,  
 992 68(2).

993 Grabowski, R.C., Droppo, I.G. and Wharton, G., 2011. Erodibility of cohesive sediment: The  
 994 importance of sediment properties. *Earth-Science Reviews*, 105(3-4), 101-120.

995 Grinsted, A., Moore, J. C., and Jevrejeva, S., 2013, Projected Atlantic hurricane surge threat  
 996 from rising temperatures: Proceedings of the National Academy of Sciences of the  
 997 United States of America, v. 110, no. 14, p. 5369-5373.

998 Haddad, J., Lawler, S., and Ferreira, C. M., 2016, Assessing the relevance of wetlands for storm  
 999 surge protection: a coupled hydrodynamic and geospatial framework: *Natural Hazards*,  
 1000 v. 80, no. 2, p. 839-861.

1001 Haggart, B.A. (1988) The stratigraphy, depositional environment and dating of a possible tidal  
 1002 surge deposit in the Beaulieu Firth area, northeast Scotland. *Palaeogeography,*  
 1003 *Palaeoclimatology, Palaeoecology* 66(3), 215-230.

1004 Hawkes, A.D., Horton, B.P. (2012) Sedimentary record of storm deposits from Hurricane Ike,  
 1005 Galveston and San Luis Islands, Texas, *Geomorphology*  
 1006 doi:10.1016/j.geomorph.2012.05.017

1007 Heuner, M., Silinski, A., Schoelynck, J., Bouma, T.J., Puijalon, S., Troch, P., Fuchs, E.,  
 1008 Schroder, B., Schroder, U., Meire, P., Temmerman, S., 2015. Ecosystem  
 1009 engineering by plants on wave-exposed intertidal flats is governed by  
 1010 relationships between effect and response traits. *PLoS One* 10, e0138086,  
 1011 <http://dx.doi.org/10.1371/journal.pone.0138086>  
 1012 <http://dx.doi.org/10.1371/journal.pone.0138086>.

1013 Hartnall, T.J., 1984. Salt-marsh vegetation and micro-relief development on the New Marsh at  
 1014 Gibraltar Point, Lincolnshire. In: Clark, M.W. (Ed.), *Coastal Research: UK*  
 1015 *Perspectives*. Cambridge University Press, Cambridge, pp. 37-58, 131 pp.

1016 Hendrickson, J.C., 1997. *Coastal wetland response to rising sea-level: quantification of short-*  
 1017 *and long-term accretion and subsidence, northeastern Gulf of Mexico*.

1018 Horton, B.P., Rossi, V. and Hawkes, A.D., 2009. The sedimentary record of the 2005 hurricane  
 1019 season from the Mississippi and Alabama coastlines. *Quaternary International*, 195(1-  
 1020 2), 15-30.

1021 Hu, K. L., Chen, Q., and Wang, H. Q., 2015, A numerical study of vegetation impact on  
 1022 reducing storm surge by wetlands in a semi-enclosed estuary: *Coastal Engineering*, v.  
 1023 95, p. 66-76.

1024 Jiang, P., Yu, Z., Gautam, M.R., Yuan, F. and Acharya, K., 2016. Changes of storm properties  
 1025 in the United States: observations and multimodel ensemble projections. *Global and*  
 1026 *Planetary Change*, 142, pp.41-52.

1027 Kearney, M.S., Grace, R.E. and Stevenson, J.C., 1988. Marsh Loss in Nanticoke Estuary,  
 1028 Chesapeake Bay. *Geographical Review*, 78(2), 205-220.

1029 Kirwan, M.L., Guntenspergen, G.R., D'Alpaos, A., Morris, J.T., Mudd, S.M. and Temmerman,  
 1030 S., 2010a. Limits on the adaptability of coastal marshes to rising sea level. *Geophysical*  
 1031 *Research Letters*, 37(23), L23401.

1032 Kirwan, M.L. and Guntenspergen, G.R., 2010b. Influence of tidal range on the stability of  
 1033 coastal marshland. *Geophysical Research Letters*, 115(F2), F02009.

1034 Kirwan, M.L., Murray, A.B., Donnelly, J.P. and Corbett, D.R., 2011. Rapid wetland expansion  
 1035 during European settlement and its implication for marsh survival under modern  
 1036 sediment delivery rates. *Geology*, 39(5), pp.507-510.

1037 Kirwan, M.L. and Mudd, S.M., 2012. Response of salt-marsh carbon accumulation to climate  
 1038 change. *Nature*, 489(7417), p.550.

1039 Kirwan, M.L., Temmerman, S., Skeehean, E.E., Guntenspergen, G.R. and Fagherazzi, S., 2016.  
 1040 Overestimation of marsh vulnerability to sea level rise. *Nature Clim. Change*, 6(3), 253-  
 1041 260.

1042 Kolker, A.S., Goodbred Jr, S.L., Hameed, S. and Cochran, J.K., 2009. High-resolution records  
 1043 of the response of coastal wetland systems to long-term and short-term sea-level  
 1044 variability. *Estuarine, Coastal and Shelf Science*, 84(4), 493-508.

1045 Kobayashi, N., Raichle, A.W. and Asano, T., 1993. Wave attenuation by vegetation. *Journal*  
 1046 *of waterway, port, coastal, and ocean engineering*, 119(1), pp.30-48.

1047 Koch, E.W. and Gust, G., 1999. Water flow in tide-and wave-dominated beds of the seagrass  
 1048 *Thalassia testudinum*. *Marine Ecology Progress Series*, 184, pp.63-72.

1049 Krauss, K. W., Doyle, T. W., Doyle, T. J., Swarzenski, C. M., From, A. S., Day, R. H., and  
1050 Conner, W. H., 2009, Water level observations in mangrove swamps during two  
1051 hurricanes in Florida: *Wetlands*, v. 29, no. 1, p. 142-149.

1052 Lawler, S., Haddad, J., and Ferreira, C. M., 2016, Sensitivity considerations and the impact of  
1053 spatial scaling for storm surge modeling in wetlands of the Mid-Atlantic region: *Ocean*  
1054 *& Coastal Management*, v. 134, p. 226-238.

1055 Lee, M., 2001. Coastal defence and the Habitats Directive: predictions of habitat change in  
1056 England and Wales. *The Geographical Journal*, 167(1), pp.39-56.

1057 Le Hir, P., Monbet, Y. and Orvain, F., 2007. Sediment erodability in sediment transport  
1058 modelling: Can we account for biota effects? *Continental Shelf Research*, 27(8), 1116-  
1059 1142.

1060 Leonard, L.A., Hine, A.C., Luther, M.E., Stumpf, R.P. and Wright, E.E., 1995. Sediment  
1061 transport processes in a west-central Florida open marine marsh tidal creek; the role of  
1062 tides and extra-tropical storms. *Estuarine, Coastal and Shelf Science*, 41(2), pp.225-  
1063 248.

1064 Le Hir, P., Roberts, W., Cazaillet, O., Christie, M., Bassoullet, P., Bacher, C., 2000.  
1065 Characterization of intertidal flat hydrodynamics. *Continental Shelf Research* 20,  
1066 1433-1459.

1067 Leonardi, N. and Fagherazzi, S., 2014. How waves shape salt marshes. *Geology*, 42(10),  
1068 pp.887-890.

1069 Leonardi, N. and Fagherazzi, S., 2015. Effect of local variability in erosional resistance on l  
1070 arge-scale morphodynamic response of salt marshes to wind waves and extreme events.  
1071 *Geophysical Research Letters*, 42(14), pp.5872-5879.

1072 Leonardi, N., Ganju, N.K. and Fagherazzi, S., 2016a. A linear relationship between wave  
 1073 power and erosion determines salt-marsh resilience to violent storms and hurricanes.  
 1074 *Proceedings of the National Academy of Sciences*, 113(1), pp.64-68.

1075 Leonardi, N., Defne, Z., Ganju, N.K. and Fagherazzi, S., 2016b. Salt marsh erosion rates and  
 1076 boundary features in a shallow Bay. *Journal of Geophysical Research: Earth Surface*,  
 1077 121(10), pp.1861-1875.

1078 Liu, K., Fearn, M.L. (2000) Reconstruction of prehistoric landfall frequencies of catastrophic  
 1079 hurricanes in northwestern Florida from lake sediment records. *Quaternary Research*  
 1080 54, 238–245.

1081 Liu, H. Q., Zhang, K. Q., Li, Y. P., and Xie, L., 2013, Numerical study of the sensitivity of  
 1082 mangroves in reducing storm surge and flooding to hurricane characteristics in southern  
 1083 Florida: *Continental Shelf Research*, v. 64, p. 51-65.

1084 Loder, N. M., Irish, J. L., Cialone, M. A., and Wamsley, T. V., 2009, Sensitivity of hurricane  
 1085 surge to morphological parameters of coastal wetlands: *Estuarine Coastal and Shelf*  
 1086 *Science*, v. 84, no. 4, p. 625-636.

1087 Lovelace, J. K., 1994, Storm-tide elevations produced by Hurricane Andrew along the  
 1088 Louisiana coast, August 25-27, 1992, Open File Report 94-371.

1089 Luhar, M., Nepf, H.M., 2016. Wave-induced dynamics of flexible blades. *J. Fluid Struct.* 61,  
 1090 20–41.

1091 Luhar, M., Coutu, S., Infantes, E., Fox, S. and Nepf, H., 2010. Wave-induced velocities  
 1092 inside a model seagrass bed. *Journal of Geophysical Research: Oceans*, 115(C12).  
 1093

1094 Ma, Z., Ysebaert, T., van der Wal, D., de Jong, D.J., Li, X. and Herman, P.M.J., 2014. Long-  
 1095 term salt marsh vertical accretion in a tidal bay with reduced sediment supply.  
 1096 *Estuarine, Coastal and Shelf Science*, 146, 14-23.

1097 Marani, M., d'Alpaos, A., Lanzoni, S. and Santalucia, M., 2011. Understanding and  
 1098 predicting wave erosion of marsh edges. *Geophysical Research Letters*, 38(21).  
 1099

1100 Mariotti, G. and Fagherazzi, S., 2010. A numerical model for the coupled long-term evolution  
 1101 of salt marshes and tidal flats. *Journal of Geophysical Research: Earth*  
 1102 *Surface*, 115(F1).

1103 Mariotti, G. and Fagherazzi, S., 2013. Critical width of tidal flats triggers marsh collapse in the  
 1104 absence of sea-level rise. *Proceedings of the national Academy of Sciences*, 110(14),  
 1105 pp.5353-5356.

1106 Mariotti, G. and Carr, J., 2014. Dual role of salt marsh retreat: Long-term loss and short-term  
 1107 resilience. *Water Resources Research*, 50(4), pp.2963-2974.

1108 Mariotti, G., 2016. Revisiting salt marsh resilience to sea level rise: Are ponds responsible for  
 1109 permanent land loss?. *Journal of Geophysical Research: Earth Surface*, 121(7),  
 1110 pp.1391-1407.

1111 Marsooli, R., Orton, P. M., Georgas, N., and Blumberg, A. F., 2016, Three-dimensional  
 1112 hydrodynamic modeling of coastal flood mitigation by wetlands: Coastal Engineering,  
 1113 v. 111, p. 83-94.

1114 McGee, B. D., Goree, B. B., Tollet, R. W., Woodward, B. K., and Kress, W. H., 2006,  
 1115 Hurricane Rita surge data, Southwestern Louisiana and Southeastern Texas,  
 1116 September-November 2005: US Geological Survey.

1117 McCave, I.N., 1987. Fine sediment sources and sinks around the East Anglian Coast (UK).  
 1118 *Journal of the Geological Society*, 144(1), 149-152.

1119 McKee, K.L. and Cherry, J.A., 2009. Hurricane Katrina sediment slowed elevation loss in  
 1120 subsiding brackish marshes of the Mississippi River delta. *Wetlands*, 29(1), 2-15.

1121 Meire, P., Dauwe, W., Maris, T., Peeters, P., Coen, L., Deschamps, M., Rutten, J., and  
 1122 Temmerman, S., 2014, Sigma plan proves efficiency: ECSA Bulletin, v. 62, p. 19-23.

1123 Mendez, F. J., and I. J. Losada (2004), An empirical model to estimate the propagation of  
 1124 random breaking and non-breaking waves over vegetation fields, *Coastal Eng.* , 51,  
 1125 103–118, doi:10.1016/j.coastaleng.2003.11.003.

1126 Miselis, J.L., Andrews, B.D., Nicholson, R.S., Defne, Z., Ganju, N.K. and Navoy, A., 2016.  
 1127 Evolution of mid-Atlantic coastal and back-barrier estuary environments in response to  
 1128 a hurricane: Implications for barrier-estuary connectivity. *Estuaries and Coasts*, 39(4),  
 1129 pp.916-934.

1130 Möller, I., T. Spencer, J.R. French, D.J. Leggett, and M. Dixon (1999), Wave transformation  
 1131 over salt marshes: a field and numerical modelling study from North Norfolk,  
 1132 England. *Estuarine Coastal Shelf Science* 49: 411–426.

1133 Moller I. 2006. Quantifying saltmarsh vegetation and its effect on wave height dissipation:  
 1134 results from a UK east coast saltmarsh. *Estuar. Coast. Shelf Sci.* 69:337–51.

1135 Morton, R.A., Sallenger, A.H., 2003. Morphological impacts of extreme storms on sandy  
 1136 beaches and barriers. *Journal of Coastal Research* 19, 560–573.

1137 Morton, R.A., Gelfenbaum, G., Jaffe, B.E. (2007) Physical criteria for distinguishing sandy  
 1138 tsunami and storm deposits using modern examples. *Sedimentary Geology* 200 (3–4),  
 1139 184-207

1140 Morton, R.A. and Barras, J.A., 2011. Hurricane impacts on coastal wetlands: A half-century  
 1141 record of storm-generated features from southern Louisiana. *Journal of Coastal*  
 1142 *Research*, 27(6A), pp.27-43.

1143 Mudd, S.M., Howell, S.M. and Morris, J.T., 2009. Impact of dynamic feedbacks between  
 1144 sedimentation, sea-level rise, and biomass production on near-surface marsh



1145 stratigraphy and carbon accumulation. *Estuarine, Coastal and Shelf Science*, 82(3),  
 1146 pp.377-389.

1147

1148 Mullarney, J.C., Henderson, S.M. (2010), Wave-forced motion of submerged single-stem  
 1149 vegetation, *J. Geophys. Res.*, 115, C12061, doi:10.1029/2010JC006448.

1150 Murray, A.L. and Spencer, T., 1997. On the wisdom of calculating annual material budgets in  
 1151 tidal wetlands. *Marine Ecology Progress Series*, 150, pp.207-216.

1152 Nicholls, R.J., Dredge, A. and Wilson, T., 2000. Shoreline change and fine-grained sediment  
 1153 input: Isle of Sheppey Coast, Thames Estuary, UK. Geological Society, London,  
 1154 Special Publications, 175(1), 305-315.

1155 Neumeier, U., Ciavola, P., 2004. Flow resistance and associated sedimentary processes in a  
 1156 *Spartina maritima* salt-marsh. *J. Coast. Res.* 20, 435–447.

1157 Paul, M., Bouma, T.J., Amos, C.L., 2012. Wave attenuation by submerged vegetation:  
 1158 combining the effect of organism traits and tidal current. *Marine Ecology-Progress*  
 1159 *Series* 444, 31–41.

1160 Paul, M., Rupprecht, F., Möller, I., Bouma, T.J., Spencer, T., Kudella, M., Wolters, G., van  
 1161 Wesenbeeck, B.K., Jensen, K., Miranda-Lange, M., Schimmels, S., 2016. Plant  
 1162 stiffness and biomass as drivers for drag forces under extreme wave loading: a flume  
 1163 study on mimics. *Coast. Eng.* 117, 70–78.

1164

1165 Pachauri, R.K., Allen, M.R., Barros, V.R., Broome, J., Cramer, W., Christ, R., Church, J.A.,  
 1166 Clarke, L., Dahe, Q., Dasgupta, P. and Dubash, N.K., 2014. *Climate change 2014:*  
 1167 *synthesis report. Contribution of Working Groups I, II and III to the fifth assessment*  
 1168 *report of the Intergovernmental Panel on Climate Change* (p. 151). IPCC.

1169 Pendleton, L., Donato, D.C., Murray, B.C., Crooks, S., Jenkins, W.A., Sifleet, S., Craft, C.,  
 1170 Fourqurean, J.W., Kauffman, J.B., Marbà, N. and Megonigal, P., 2012. Estimating  
 1171 global “blue carbon” emissions from conversion and degradation of vegetated coastal  
 1172 ecosystems. *PloS one*, 7(9), p.e43542.

1173 Peralta, G., van Duren, L.A., Morris, E.P., Bouma, T.J., 2008. Consequences of shoot density  
 1174 and stiffness for ecosystem engineering by benthic macrophytes in flow dominated  
 1175 areas: a hydrodynamic flume study. *Mar. Ecol.—Prog. Ser.* 368, 103–115.

1176 Pethick, J.S., 1992. Saltmarsh geomorphology. In: Allen, J.R.L., Pye, K. (Eds.), *Saltmarshes.*  
 1177 *Morphodynamics, Conservation and Engineering Significance.* Cambridge University  
 1178 Press, Cambridge, pp. 41-62.

1179 Puijalon, S., Bouma, T.J., Douady, C.J., van Groenendael, J., Anten, N.P.R., Martel, E.,  
 1180 Bornette, G., 2011. Plant resistance to mechanical stress: evidence of an  
 1181 avoidance-tolerance trade-off. *New Phytologist*, 191, pp. 1141–1149.

1182 *Rahman, R., Plater, A. J., Nolan, P. J., Mauz, B., Appleby, P. G. (2013). Potential health risks*  
 1183 *from radioactive contamination of saltmarshes in NW England. Journal of*  
 1184 *Environmental Radioactivity* 119, 55-62.

1185 Redfield, A.C., 1972. Development of a New England salt marsh. *Ecological*  
 1186 *monographs*, 42(2), pp.201-237.

1187 Reed, D.J., 1989. Patterns of sediment deposition in subsiding coastal salt marshes, Terrebonne  
 1188 Bay, Louisiana: the role of winter storms. *Estuaries and Coasts*, 12(4), pp.222-227.

1189 Reed, D.J., 2002. Understanding tidal marsh sedimentation in the Sacramento-San Joaquin  
 1190 Delta. *California. Journal of Coastal Research Special*, (36), pp.605-611.

1191 Resio, D. T., and Westerink, J. J., 2008, Modelling the physics of storm surges: *Physics Today*,  
 1192 v. 61, no. 9, p. 33-38.

1193 Riffe, K. C., S. M. Henderson, and J. C. Mullarney (2011), Wave dissipation by flexible  
 1194 vegetation, *Geophysical Research Letters*, 38,p. L18607,  
 1195 doi:10.1029/2011GL048773.

1196 Rosencranz, J.A., Ganju, N.K., Ambrose, R.F., Brosnahan, S.M., Dickhudt, P.J.,  
 1197 Guntenspergen, G.R., MacDonald, G.M., Takekawa, J.Y. and Thorne, K.M., 2016.  
 1198 Balanced Sediment Fluxes in Southern California's Mediterranean-Climate Zone Salt  
 1199 Marshes. *Estuaries and Coasts*, 39(4), pp.1035-1049.

1200 Rupprecht, F., Moller, I., Evans, B., Spencer, T., and Jensen, K., 2015, Biophysical properties  
 1201 of salt marsh canopies Quantifying plant stem flexibility and above ground biomass:  
 1202 *Coastal Engineering*, v. 100, p. 48-57.

1203 Rupprecht, F., Möller, I., Paul, M., Kudella, M., Spencer, T., van Wesenbeeck, B. K., ... &  
 1204 Schimmels, S. (2017). Vegetation-wave interactions in salt marshes under storm surge  
 1205 conditions. *Ecological Engineering*, 100, 301-315.

1206

1207 Schepers, L., Kirwan, M. L., Guntenspergen, G. R., and Temmerman, S., 2017, Spatio-  
 1208 temporal development of vegetation die-off in a submerging coastal marsh: *Limnology*  
 1209 and *Oceanography*, v. 62, p. 137-150.

1210 Schuerch, M., Rapaglia, J., Liebetrau, V., Vafeidis, A. and Reise, K., 2012. Salt Marsh  
 1211 Accretion and Storm Tide Variation: an Example from a Barrier Island in the North  
 1212 Sea. *Estuaries and Coasts*, 35(2), 486-500.

1213 Schuerch, M., Vafeidis, A., Slawig, T. and Temmerman, S., 2013. Modeling the influence of  
 1214 changing storm patterns on the ability of a salt marsh to keep pace with sea level rise.  
 1215 *Journal of Geophysical Research: Earth Surface*, 118(1), 84-96.

1216 Schuerch, M., Dolch, T., Reise, K. and Vafeidis, A.T., 2014. Unravelling interactions between  
 1217 salt marsh evolution and sedimentary processes in the Wadden Sea (southeastern North  
 1218 Sea). *Progress in Physical Geography*, 38(6), 691-715.

1219 Schuerch, M., Scholten, J., Carretero, S., García-Rodríguez, F., Kumbier, K., Baechtger, M.  
 1220 and Liebetrau, V., 2016. The effect of long-term and decadal climate and hydrology  
 1221 variations on estuarine marsh dynamics: An identifying case study from the Río de la  
 1222 Plata. *Geomorphology*, 269, 122-132.

1223 Schwimmer, R.A., 2001. Rates and processes of marsh shoreline erosion in Rehoboth Bay,  
 1224 Delaware, USA. *Journal of Coastal Research*, pp.672-683.

1225 Scileppi, E. and Donnelly, J.P., 2007. Sedimentary evidence of hurricane strikes in western  
 1226 Long Island, New York. *Geochemistry, Geophysics, Geosystems*, 8(6).

1227 *Scileppi, E., Donnelly, J.P. (2007) Sedimentary evidence of hurricane strikes in western Long*  
 1228 *Island, New York. Geochemistry, Geophysics and Geosystems* 8(6), Q06011,  
 1229 doi:10.1029/2006GC001463

1230 Sheng, Y. P., Lapetina, A., and Ma, G. F., 2012, The reduction of storm surge by vegetation  
 1231 canopies: Three-dimensional simulations: *Geophysical Research Letters*, v. 39.  
 1232  
 1233

1234 Silinski, A., Heuner, M., Schoelynck, J., Puijalon, S., Schroder, U., Fuchs, E., Troch, P.,  
 1235 Bouma, T.J., Meire, P., Temmerman, S., 2015. Effects of wind waves versus ship  
 1236 waves on tidal marsh plants: a flume study on different life stages of *Scirpus maritimus*.  
 1237 *PLoS One* 10, e0118687,  
 1238 [http://dx.doi.org/10.1371/journal.ponehttp://dx.doi.org/10.1371/journal.pone.](http://dx.doi.org/10.1371/journal.ponehttp://dx.doi.org/10.1371/journal.pone.0118687)  
 1239 0118687.  
 1240

1241 Smith, D.E., Shi, S., Cullingford, R.A., Dawson, A.G., Dawson, S., Firth, C.R., Foster, I.D.L.,  
1242 Fretwell, P.T., Haggart, B.A. Holloway, L.K., Long, D. (2004) The Holocene Storegga  
1243 Slide tsunami in the United Kingdom Quaternary Science Reviews 23(23–24), 2291-  
1244 2321.

1245 Smolders, S., Plancke, Y., Ides, S., Meire, P., and Temmerman, S., 2015, Role of intertidal  
1246 wetlands for tidal and storm tide attenuation along a confined estuary: a model study:  
1247 Natural Hazards and Earth System Sciences, v. 15, p. 1659–1675.

1248 Sobel, A.H., Held, I.M. and Bretherton, C.S., 2002. The ENSO signal in tropical tropospheric  
1249 temperature. *Journal of Climate*, 15(18), pp.2702-2706.

1250 Solomon, S. ed., 2007. *Climate change 2007-the physical science basis: Working group I*  
1251 *contribution to the fourth assessment report of the IPCC* (Vol. 4). Cambridge  
1252 University Press.

1253 Spencer, T., Brooks, S.M., Evans, B.R., Tempest, J.A. and Möller, I., 2015. Southern North  
1254 Sea storm surge event of 5 December 2013: water levels, waves and coastal impacts.  
1255 *Earth-Science Reviews*, 146, pp.120-145.

1256 Spencer, T., Moller, I., Rupprecht, F., Bouma, T.J., van Wesenbeeck, B.K., Kudella, M., Paul,  
1257 M., Jensen, K., Wolters, G., Miranda-Lange, M., Schimmels, S., 2016. Salt marsh  
1258 surface survives true-to-scale simulated storm surges. *Earth Surface Processes*  
1259 *and Landforms*. DOI:10.1002/esp.3867.

1260 Spencer, T., Schuerch, M., Nicholls, R.J., Hinkel, J., Lincke, D., Vafeidis, A.T., Reef, R.,  
1261 McFadden, L. and Brown, S., 2016. Global coastal wetland change under sea-level rise  
1262 and related stresses: The DIVA Wetland Change Model. *Global and Planetary Change*,  
1263 139, 15-30.

1264 Stark, J., Plancke, Y., Ides, S., Meire, P., and Temmerman, S., 2016, Coastal flood protection  
1265 by a combined nature-based and engineering approach: modeling the effects of marsh  
1266 geometry and surrounding dikes: *Estuarine Coastal and Shelf Science*, v. 175, p. 34-45.

1267 Stark, J., Van Oyen, T., Meire, P., and Temmerman, S., 2015, Observations of tidal and storm  
1268 surge attenuation in a large tidal marsh: *Limnology and Oceanography*, v. 60, no. 4, p.  
1269 1371-1381.

1270 Stevenson, J.C., Kearney, M.S. and Pendleton, E.C., 1985. Sedimentation and erosion in a  
1271 Chesapeake Bay brackish marsh system. *Marine Geology*, 67(3-4), pp.213-235.

1272 Stumpf, R.P. (1983) The process of sedimentation on the surface of a salt marsh. *Estuarine,*  
1273 *Coastal and Shelf Science* 17(5), 495-508.

1274 Sutton-Grier, A. E., Wowk, K., and Bamford, H., 2015, Future of our coasts: The potential for  
1275 natural and hybrid infrastructure to enhance the resilience of our coastal communities,  
1276 economies and ecosystems: *Environmental Science & Policy*, v. 51, p. 137-148.

1277 Syvitski, J.P.M., Vörösmarty, C.J., Kettner, A.J. and Green, P., 2005. Impact of Humans on  
1278 the Flux of Terrestrial Sediment to the Global Coastal Ocean. *Science*, 308(5720), 376-  
1279 380.

1280 Temmerman, S., Govers, G., Meire, P. and Wartel, S., 2003. Modelling long-term tidal marsh  
1281 growth under changing tidal conditions and suspended sediment concentrations,  
1282 Scheldt estuary, Belgium. *Marine Geology*, 193(1-2), 151-169.

1283 Temmerman, S. et al. Ecosystem-based coastal defence in the face of global change. *Nature*  
1284 504, 79–83 (2013).

1285 Temmerman, S., De Vries, M. B., and Bouma, T. J., 2012, Coastal marsh die-off and reduced  
1286 attenuation of coastal floods: a model analysis: *Global and Planetary Change*, v. 92-93,  
1287 p. 267-274.

1288 Temmerman, S., and Kirwan, M. L., 2015, Building land with a rising sea: *Science*, v. 349, no.  
1289 6248, p. 588-589.

1290 Temmerman, S., Meire, P., Bouma, T. J., Herman, P. M. J., Ysebaert, T., and De Vriend, H. J.,  
1291 2013, Ecosystem-based coastal defence in the face of global change: *Nature*, v. 504, p.  
1292 79-83.

1293 Tonelli, M., Fagherazzi, S. and Petti, M., 2010. Modeling wave impact on salt marsh  
1294 boundaries. *Journal of Geophysical Research: Oceans*, 115(C9).

1295 Turner, R.E., Baustian, J.J., Swenson, E.M., Spicer, J.S. (2006) Wetland sedimentation from  
1296 hurricanes Katrina and Rita. *Science* 314, 449–452.

1297 Turner, R.E., Swenson, E.M., Milan, C.S., Lee J.M. (2007) Hurricane signals in salt marsh  
1298 sediments: Inorganic sources and soil volume. *Limnology and Oceanography* 52(3),  
1299 1231–1238.

1300 Tweel, A.W. and Turner, R.E., 2014. Contribution of tropical cyclones to the sediment budget  
1301 for coastal wetlands in Louisiana, USA. *Landscape Ecology*, 29(6), 1083-1094.

1302 USACE, 1963, Interim survey report, Morgan city, Louisiana and vicinity, serial no. 63: US  
1303 Army Corps of Engineers, New Orleans.

1304 Van de Koppel, J., Van der Wal, D., Bakker, J. P., and Herman, P. M. J., 2005, Self-  
1305 organisation and vegetation collapse in salt marsh ecosystems: *The American*  
1306 *Naturalist*, v. 165, no. 1, p. E1-E12.

1307 Van Wesenbeeck, B., Mulder, J. P. M., Marchand, M., Reed, D. J., De Vries, M. B., De Vriend,  
1308 H. J., and Herman, P. M. J., 2014, Damming deltas: a practice of the past? Towards  
1309 nature-based flood defenses: *Estuarine Coastal and Shelf Science*, v. 140, p. 1-6.

1310 Vecchi, G.A. and Soden, B.J., 2007. Increased tropical Atlantic wind shear in model  
1311 projections of global warming. *Geophysical Research Letters*, 34(8).

1312 Wamsley, T. V., Cialone, M. A., Smith, J. M., Atkinson, J. H., and Rosati, J. D., 2010, The  
1313 potential of wetlands in reducing storm surge: *Ocean Engineering*, v. 37, no. 1, p. 59-  
1314 68.

1315 Wamsley, T. V., Cialone, M. A., Smith, J. M., Ebersole, B. A., and Grzegorzewski, A. S., 2009,  
1316 Influence of landscape restoration and degradation on storm surge and waves in  
1317 southern Louisiana: *Natural Hazards*, v. 51, p. 207-224.

1318 Webster, P.J., Holland, G.J., Curry, J.A. and Chang, H.R., 2005. Changes in tropical cyclone  
1319 number, duration, and intensity in a warming environment. *Science*, 309(5742),  
1320 pp.1844-1846.

1321 Williams, H.F., 2009. Stratigraphy, sedimentology, and microfossil content of Hurricane Rita  
1322 storm surge deposits in southwest Louisiana. *Journal of Coastal Research*, pp.1041-  
1323 1051.

1324 Williams, H.F.L. (2011). Stratigraphic Record of Hurricanes Audrey, Rita and Ike in the  
1325 Chenier Plain of Southwest Louisiana. *Journal of Coastal Research* SI 64, 1921-1926.

1326 Woodruff, J. D., Irish, J. L., and Camargo, S. J., 2013, Coastal flooding by tropical cyclones  
1327 and sea-level rise: *Nature*, v. 504, p. 44-52.


1328 Woth, K., Weisse, R. and Von Storch, H., 2006. Climate change and North Sea storm surge  
1329 extremes: an ensemble study of storm surge extremes expected in a changed climate  
1330 projected by four different regional climate models. *Ocean Dynamics*, 56(1), pp.3-15.


1331 Yang, S.L., Shi, B.W., Bouma, T.J., Ysebaert, T., Luo, X.X., 2012. Wave attenuation at a salt  
1332 marsh margin: a case study of an exposed coast on the Yangtze Estuary. *Estuaries*  
1333 *Coasts* 35, 169– 182.


1334 Zhang, K., Douglas, B.C. and Leatherman, S.P., 2000. Twentieth-century storm activity  
1335 along the US east coast. *Journal of Climate*, 13(10), pp.1748-1761.





1336 Zhang, K. Q., Liu, H. Q., Li, Y. P., Xu, H. Z., Shen, J., Rhome, J., and Smith, T. J., 2012, The  
1337 role of mangroves in attenuating storm surges: Estuarine Coastal and Shelf Science, v.  
1338 102, p. 11-23.  
1339  
1340  
1341














A Contribution to marsh erosion of different wind categories




B Impact of extreme events on marsh shoreline shape




Fishing Bay, MD



Blackwater, MD



Dinner Creek, NJ



Reedy Creek, NJ

